

UNIVERSAL  
LIBRARY



103 325

UNIVERSAL  
LIBRARY

523.8 M38a  
Martin  
The friendly stars

68-09341

kansas city



public library

kansas city, missouri

Books will be issued only  
on presentation of library card.  
Please report lost cards and  
change of residence promptly.  
Card holders are responsible for  
all books, records, films, pictures  
or other library materials  
checked out on their cards.







# THE FRIENDLY STARS

*A New Edition of an Astronomical Classic*

Martha Evans Martin

*And*

Donald Howard Menzel



*With New Illustrations by*

CHING-SUNG YÜ

*Professor of Astronomy, Hood College*

*And a Foreword by*

DONALD HOWARD MENZEL

*and*

WILLIAM W. MORGAN

*Former Director, Yerkes Observatory*

DOVER PUBLICATIONS, INC., NEW YORK

Copyright © 1907, 1934 by Elizabeth J. Evans.  
Copyright © 1964, 1966 by Dover Publications,  
Inc.

All rights reserved under Pan American and International Copyright Conventions.

Published in Canada by General Publishing Company, Ltd., 30 Lesmill Road, Don Mills, Toronto, Ontario.

Published in the United Kingdom by Constable and Company, Ltd., 10 Orange Street, London W. C. 2.

This Dover edition, first published in 1964, is a revised and enlarged version of the work first published by Harper & Brothers in 1907.

*Library of Congress Catalog Card Number: 63-17911*

Manufactured in the United States of America

Dover Publications, Inc.  
180 Varick Street  
New York 14, N. Y.

## Foreword

This is a charming book, a book that brings us close to the starry night and other aspects of Nature that appeal to our senses. It is a Nature book in the most popular sense; it is not a book for the technical student.

And yet, *The Friendly Stars* was an important influence in the decision of two boys (living in widely separated places) to become astronomers, some forty years ago. The interest and excitement it generated were among the principal causes for our deciding to devote our lives to discovering more about the stars and the universe.

In the past we have frequently recommended *The Friendly Stars* to other young people—and have regretted that it has long been out of print—as well as in need of revision. This new, revised edition, we believe, fills a gap in the literature of elementary astronomy.

1963

DONALD H. MENZEL

*Harvard College Observatory*

WILLIAM W. MORGAN

*Yerkes Observatory*



# Acknowledgments

In revising *The Friendly Stars* I have tried, as far as possible, to retain the informal style of the original author, Martha Evans Martin. Major revision has been necessary only in the sections related to the physical characteristics of the stars: distances, velocities, temperatures, and evolutionary stages. I have also introduced the results of historical studies that I have personally made in the origin of the major constellation figures.

On the rare occasions where I differ with Mrs. Martin, I have expressed her viewpoint even when I may hold another. Our major point of disagreement concerns the relative importance of the stars and constellations; Mrs. Martin finds the star groups and constellation figures uninteresting, unimaginative, and hard to see. I believe that the constellation forms have evolved imaginatively from early times. When properly drawn, they are helpful guides to recognition of the various star groups. The figures here employed differ significantly from those given in the original edition of *The Friendly Stars*. They are identical with those I devised for *Field Guide to the Stars and Planets*.<sup>\*</sup> Many of these figures are based, in turn, on the imaginative diagrams designed by H. A. Rey, in his book *The Stars—A New Way To See Them*.<sup>†</sup> I am indebted to Mr. Rey and to Houghton Mifflin Company for permission to employ these figures in this book.

I also wish to acknowledge the excellent art work of Ching-Sung Yü. Dr. William W. Morgan, whose interest in this book encouraged the revision, not only joined me in preparation of a Foreword, but also kindly read the manuscript and made many helpful suggestions. I sincerely hope that young amateur astronomers, the world over, will find this new edition of *The Friendly Stars* interesting and stimulating.

D. H. M.

<sup>\*</sup>*Field Guide to the Stars and Planets* by Donald H. Menzel. Published by Houghton Mifflin Company. Copyright © 1963 by Donald H. Menzel.

<sup>†</sup>*The Stars—A New Way To See Them* by H. A. Rey. Published by Houghton Mifflin Company. Copyright © 1952 by H. A. Rey.



# Contents

<i>chapter</i>		<i>page</i>
	INTRODUCTION . . . . .	1
I	THE RISING AND THE SETTING OF THE STARS . . . . .	5
II	CONCERNING THE BRIGHTEST STARS . . . . .	10
III	CAPELLA . . . . .	17
IV	ARCTURUS . . . . .	20
V	SPICA . . . . .	24
VI	VEGA . . . . .	26
VII	DENEK . . . . .	30
VIII	ALTAIR . . . . .	35
IX	ANTARES . . . . .	40
X	FOMALHAUT . . . . .	45
XI	ALDEBARAN . . . . .	48
XII	ORION'S BRIGHT STARS . . . . .	52
XIII	THE HEAVENLY TWINS . . . . .	57
XIV	THE TWO DOG STARS . . . . .	61
XV	REGULUS . . . . .	66
XVI	THE NUMBER OF THE STARS . . . . .	70
XVII	THE NAMES OF THE STARS . . . . .	74
XVIII	THE LIGHT OF THE STARS . . . . .	78
XIX	THE DISTANCES OF THE STARS . . . . .	84
XX	DOUBLE STARS . . . . .	91
XXI	THE CONSTELLATIONS . . . . .	98
XXII	THE LITTLE BEAR AND POLARIS . . . . .	101
XXIII	THE GREAT BEAR AND THE SEVEN STARS OF THE DIPPER . . . . .	104
XXIV	CASSIOPEIA, CEPHEUS, AND THE DRAGON . . . . .	108
XXV	FROM THE HUNTING-DOGS TO THE ARROW . . . . .	111
XXVI	FROM THE WINGED HORSE TO THE BULL . . . . .	116
XXVII	THE MYSTERY OF THE PLEIADES . . . . .	124
XXVIII	FROM THE TWINS BACK TO THE HUNTER . . . . .	129
XXIX	CONSTELLATIONS WHOLLY OR IN PART	
	SOUTH OF THE EQUATOR . . . . .	131
XXX	INDIVIDUALITY OF THE STARS . . . . .	139
	INDEX . . . . .	143





## Illustrations

	<i>page</i>
THE BIG DIPPER . . . . .	13
RELATIVE POSITION OF THE NORTH STAR AND THE DIPPER . . . . .	14
VEGA (IN LYRA) AS SHE APPEARS IN THE EAST . . . . .	27
VEGA AND HER ATTENDANT STARS IN THE NORTHWEST . . . . .	28
TRIANGLE FORMED BY POLARIS, DENEK, AND VEGA . . . . .	31
THE NORTHERN CROSS, PART OF CYGNUS, THE SWAN . . . . .	32
ALTAIR AND ITS TWO COMPANIONS IN AQUILA . . . . .	36
DENEK, ALTAIR, AND VEGA . . . . .	37
THE DOLPHIN (DELPHINUS), OR JOB'S COFFIN . . . . .	38
SCORPIUS . . . . .	41
FOMALHAUT . . . . .	46
THE HYADES . . . . .	49
ORION . . . . .	53
THE SICKLE . . . . .	67
THE LITTLE DIPPER . . . . .	103
THE SEVEN STARS OF THE BIG DIPPER . . . . .	105
CASSIOPELA . . . . .	109
HERCULES . . . . .	113
ANDROMEDA, THE SQUARE OF PEGASUS, AND THE SEGMENT OF PERSEUS . . . . .	118
THE PLEIADES . . . . .	125
LEO OR THE LION . . . . .	130
LIBRA OR THE SCALES . . . . .	132
SAGITTARIUS OR THE MILK DIPPER . . . . .	133
PLATE I—NORTH CIRCUMPOLAR STAR CHARTS . . . . .	<i>inside back cover</i>
PLATE II—EQUATORIAL STAR CHARTS . . . . .	<i>inside back cover</i>



## Introduction

THE chief aim of this book is to share with others the pleasure that the writer has had in what may be called a relation of personal friendship with the stars. In this relation one knows the more conspicuous stars by name and at a glance; and is able to greet them as pleasant acquaintances when they return year after year in the due seasons, or each evening as they pass over their prescribed paths serene and stately, or dancing and twinkling, according to their several habits.

One has a fine sense of companionship with the stars when he has secured this kind of acquaintance with them—when on looking out of the window at any hour of the night he can see a familiar face twinkling at him as if in friendly recognition of the fact that he must know it is due at that hour and is expecting to see it; or when, on a cold midnight in late February, before the trees and birds have announced the spring-time, he sees a bright, bluish, scintillating point just pushing up over the eastern horizon and knows that Vega has come to grace the skies again and that spring will surely come with her.

Such a feeling for the stars is not induced by exciting wonder at the expanse and mystery of the heavens, or by burdening and oppressing the mind with the vastness that seems beyond all compassing in thought, but by showing how the stars, like the flowers and the trees, are but parts of the visible beauty of nature which have their share in making "the perfect whole."

Plants and birds come in their turn

As the revolving seasons rise  
Above the tree-tops star by star,

and the steady advance of the changing season gets a definiteness and an interest to one otherwise impossible when he has learned to associate the visible signs of the progress of the year

as they appear in the skies as well as on the earth. He will then associate the blooming of the blood-root and the first warble of the bluebirds with the eastern splendor of Arcturus and the blooming of the maples. When he watches in the daytime for the first blue violets he will look the same evening for the blue twinkling face of Vega. He will know that the juncos and Sirius leave us at about the same time in the spring; that when the golden-rod and the wild asters are blooming it is time to look for Fomalhaut and that Antares is about to go; and a creaking, frosty night will make him involuntarily turn his eyes up to mighty Orion striding across the southern skies.

Such things as these, which are so dear to the heart of the lover of nature, are not learned as one learns a lesson, or by the being told, but by coming into a personal relation, a relation very easily established by one who really desires it, with the flowers and trees and birds and stars, knowing them all in their seasons and their associations.

For such knowledge as this no technical information is necessary, though it may be followed by a desire for technical knowledge. The two are quite different things. But one can come to no real appreciation of anything in nature without some knowledge of particular objects. The acquirement of it is just as important and just as easy with the stars as with the flowers. One may revel in the beauty of a whole field of grass and flowers, but his heart gives a leap when he sees among them the face of some flower that he knows and likes, and he cries, "There's a lily," or a gentian, or whatever it may be. Birds may flit around us as we sit in the woods, and we note them as black, yellow, pretty, or whatever, and we think we are getting all the pleasure we can from them; but interest quickens when one comes that we can name, and it at once has an individuality and an importance that none of the rest have.

So it is with the stars: a starry night is beautiful and we gaze at it and enjoy it and do not care to know more about it in detail. But if by chance we come to know by name one bright star, it immediately separates itself from all the others and becomes an individual. If we enlarge our acquaintance in the skies, the whole aspect of the heavens is changed, and, instead of a brilliant assembly of impersonal points of light, we see

a host of individuals that we know as bright Capella, sombre Betelgeuse, and others.

And this satisfaction we may secure without troubling about meridians and ecliptics, or right ascension and declination, or any other of the scientific trappings of the stars, just as we have it with the flowers without attending to sepals and petals, or pistils and anthers, and with the birds without any thought of anatomical classification. The only thing one needs to do in order to have such an acquaintance with the stars is to look for them: and this book aims to show how and when and where to look for them in the easiest possible way.

Any especially scientific information will be omitted, the desire being to make the stars as interesting in themselves as possible and to show how simple a matter it is to know in an intelligent but untechnical way the choicest individuals among them, the leading facts concerning them, their bearing in practical life, and all that is individually beautiful and important about them. This, it is believed, will yield a pleasure such that any one who has once experienced it will receive a very large reward for the small labor entailed.

It is by no means a dry and difficult thing to acquire any particular knowledge of the stars. One, if rightly directed, will find on pursuing the subject only just a little way that, while giving himself next to no trouble about it, he can soon come to know the more notable ones by name and to have a fair fund of practical information about them. Many of the stars, which now all look so much alike to him, will have acquired such an individuality that he can recognize them at a glance in whatever position in the skies he may see them. And he will find, too, that the pleasure one never fails to have in the spectacle of the whole sky is multiplied many times for him as soon as he knows something of the individuals that make up this splendid panorama.

It is an interesting fact that the stars as individuals figure a great deal more prominently in old than in modern literature, and the writers of the older literature show a more personal acquaintance with them than the writers of the later literature show. This intimate knowledge of the stars really preceded the science of astronomy. The stars were better known even to people at large before there was any such science than they are

known now, and there is only too much justice in Emerson's reproach that now, in these days of nautical almanacs, "the man in the street does not know a star in the sky. The solstice he does not observe, the equinox he knows as little; and the whole bright calendar of the year is without a dial in his mind."

The explanation of this difference is that people in earlier times, being more out of doors than the people of the present day, were more observant of out-door objects and took more interest in them. For it is, after all, mainly a matter of observation; and for learning to know the leading stars and getting the interest that attends an acquaintance with them, little more is needful than simply looking at them. It is looking at them, too, with only the naked eye; for all the stars that attract special notice and have individual names were noticed and so named long before the invention of the telescope; and the principal constellations were traced and named by simple shepherds who tended their flocks at night in the open fields and had nothing to aid them but their own eyes and fancy.

# I

## The Rising and the Setting of the Stars

THE most favorable time for identifying a star is when it is rising, or has just risen, especially if its rising takes place anywhere between twilight and midnight, the portion of the day most likely to be convenient for observers. It is then in the eastern part of the sky, where the afterglow of the sunset does not dim its luster, and it is far enough down towards the horizon for one to look at it face to face in a comfortable attitude. Moreover, at rising, a star is more nearly by itself, more separated from neighboring stars, and is, therefore, much more easily recognized than when one has to pick it out from among the host of other stars already scattered over the sky and in as plain view as it. For near the horizon the atmosphere obliterates the multitude of faint stars that are plainly visible higher up in the sky.

It is true that, owing to the density of the atmosphere there, all stars twinkle more, and hence shine with less steadiness, when they are near the horizon than when they are higher up in the sky, and most stars are not to be seen at all until they have actually risen some distance above the horizon. Then there is to be allowed for, the interference of buildings, trees, and other objects in the landscape; but, even in spite of these interferences, most any star comes well into view before it has risen so high that it cannot be easily separated from the others and looked at with comfort. One has the advantage given by the comparative isolation of a rising star whether it rises from below the horizon or from behind the trees and houses.

If one were observing a star through a telescope and for other purposes than that of identification, he should choose, as nearly as possible, a time when the star is in the middle of its course from east to west, or as nearly overhead as it ever reaches. There

the atmosphere will be clearer and, in consequence, the star will shine with a steadier light and offer less opportunity for optical illusions. But this means either a neck-breaking attitude or some special preparations for a comfortable reclining posture, either of which is more or less discouraging to one who has not already become pretty familiar with the stars. When he has come to know them, there is no greater delight than to lie stretched on a cot in the open with the whole expanse of the heavens above him, and watch them as they rise and pass and go again. This is the stage of acquaintance that binds one forever in friendship with the stars. But a speaking acquaintance may be maintained that will give much pleasure with lighter demands upon one.

The rising of the stars is the time of recognition and greeting. But it is no less important to speed the parting guest; and the next most satisfactory time to watch a star is when it has reached the western sky and is about to leave us until it comes again to the eastern door on its never-ceasing round. Here, while the star remains the same in its individual attributes, all its setting is the reverse of what it was when the star first appeared above the eastern horizon. The parallelogram of twinkling stars that trailed behind Vega as she rose in the east now waves above her as she sinks below the line of vision in the west. The Northern Cross, which heaved itself into sight in the east almost wrong-side up, with Deneb weighting the head of it, has, twenty hours later, practically righted itself and is lost to view in the glowing west with Deneb dancing on the tip of its upright. So complete identification must include the recognition of a star both in the east and in the west.

In the north the stars that constantly circle around the pole are always visible to us. They complete their small round once in twenty-four hours; but their path lies entirely above the horizon, so they never rise or set. Within this circle are the Pole Star and the stars of the Great Dipper, which will be illustrated in the next chapter; but it includes none of the brightest stars.

Next comes a line of stars describing a larger circle around the pole, one that takes them below the horizon for a longer or shorter time according to their distance from the pivotal point, the pole. These stars rise in the northeast, make a long, slow sweep of the sky, and set in the northwest, finishing their circle



below the horizon and rising not long after they set. Among these are some of the brightest stars.

Then follows a belt of stars that rise somewhere near the exact eastern point of the horizon, cross the heavens to the western point in about twelve hours, and remain below the horizon another twelve hours.

And, finally, there are the stars in the southern section of the sky, which rise south of east and set in the southwest, never rising very high in the heavens, the portion of their circle, or course, over which we can see them pass being smaller and smaller as that course lies farther and farther south of us. Among these southern stars are Sirius, the star that outshines all the others, and Fomalhaut, which hides from us for sixteen hours out of each twenty-four.

All the stars that rise south of the exact eastern point on the horizon are called southern stars; for the sky is divided into a northern and a southern hemisphere by an equator running half-way between the north and the south pole, just as the earth is divided. The northernmost stars of the southern hemisphere are visible to us just as the southernmost stars of the northern hemisphere are visible to people living in southern latitudes. These distinctions are of no great importance, but it is well to understand that, while we cannot see the southern stars as a body, yet some of them are visible to us and still are spoken of as southern stars. Three of the brightest stars that come within our view, Rigel, Antares, and Spica, are southern stars, in addition to Sirius and Fomalhaut, which have already been mentioned.

Unlike the sun and moon and the planets, each star rises always at practically the same point in the heavens. It comes back to the starting-point each day four minutes earlier than the day before, which makes it rise about an hour earlier each fortnight, and two hours earlier each month; so that in twelve months it comes back twenty-four hours earlier. This brings it back to the same rising point at almost the same second once a year.

These changes from day to day in the time of the rising and the setting of the stars are regular and definite; and so it is an easy matter, after one has once learned to recognize a star, either by its appearance or its surroundings, to know where to

look for it at any given date. If a star, for instance, is due to rise at six o'clock in the evening on the first of March, it will rise at four in the afternoon in April, and at eight in the evening in February; at two o'clock in the afternoon in May, and at ten in the evening in January; at noon in June, and at midnight in December; and so on throughout the year. Hence, almost any star can be seen at some time in the night for six months in the year. The other six months, if it is among the northernmost stars, it can be seen more or less each night; but if it is among those that are below the horizon many hours, it may be above it during the hours of sunshine and thus be rendered invisible to the naked eye, though even in the daytime one can usually find a bright star with a telescope.

The times of rising and setting given in this book are for about the latitude of New York City, which is nearly forty-one degrees north latitude. For any place within five degrees of this latitude on either side the difference in time of rising is so slight that for mere purposes of identification it will not matter. The time of the appearance of the stars to persons within this belt of latitude is the same all the way across the country from Maine to California. It includes the region around all the large cities of the Eastern and Middle States and most of the Western ones. Within it lie Portland, Boston, Philadelphia, Washington, Baltimore, Cincinnati, Indianapolis, Chicago, St. Paul, Detroit, Louisville, St. Louis, Denver, and San Francisco. On the southernmost edge of it is Memphis, and just a little outside of it on the north is Olympia, Washington. Los Angeles lies a little outside it on the south. Above or below this belt one must take latitude somewhat into consideration; which means only to remember that when an observer is placed farther south the stars to the north of him rise somewhat later and those to the south of him somewhat earlier, and that if he is placed north of this belt the stars to the south of him rise somewhat later and those to the north of him somewhat earlier. The rising-times of the stars as given here will answer approximately for any place in the United States.

When the rising-time of a star is referred to in these pages, it is usually given as nearly as possible in round numbers to avoid any complications. If the actual time, for instance, is 6:55, it is given as 7:00. Variations of this kind seldom extend over

more than a few minutes and they more often give a time a little later instead of earlier than the exact minute. This makes the inexactness still more immaterial, since for naked eye observation one cannot hold the stars to their scheduled minute of rising, and anywhere within half an hour is a reasonable expectation in that regard. They are there, to be sure, but variations of climate between two places and variations of atmosphere from day to day will make a considerable difference in their visibility when near the horizon. After they are well up in the sky, of course, one need not make so many concessions.

The times quoted are strictly standard times for the selected meridian. For sites to the east or west of the standard meridian the time will differ. Also for daylight saving time, remember to add one hour.

It is to be remembered also that one is rarely ever situated where he can really see the horizon, but owing to various causes sees usually a line somewhat above the true horizon.

## II

### Concerning the Brightest Stars

**A**MONG the hundreds of thousands of stars in the universe there are twenty-one that are so much brighter than the others as to make a class by themselves. They are known as stars of the first magnitude, or highest degree of brightness. These twenty-one stars are so scattered over the sky and each so dominates the section in which it is found, that if one comes to know them, there is no quarter of the heavens towards which he can turn his eye and not have a pleasant sense of familiarity with it.

Each one of these stars has a fixed place in the sky, and all but three have proper names of their own. The stars in general are not known by individual names; so it is a matter of congratulation that nearly all the bright ones are so known, and especially that this is true of all the brightest stars that can be seen in our latitude. The fixed place they have in common with all the other stars, which, when once their place is known, can always be found in the same spot. For the stars are steadfast. Spring, summer, autumn and winter each has its own bright stellar setting, and we can always count on its appearing with unchanging beauty and unfailing promptness. The moon comes and goes. She is the symbol of inconstancy. The planets wander from place to place—most of them easy enough to find, but continually changing in brightness and position. Only the stars are always the same.

The names of the twenty-one brightest stars, given in the order of their brightness, are as follows:

Sirius .....	pronounced	Sir'-i-us
Canopus .....	"	Ka-nō'-pus
Alpha Centauri .....	"	Al-fa Sen-tā'-rē
Arcturus .....	"	Ārk-tū'-rus
Vega .....	"	Vē-ga
Capella .....	"	Ka-pel'-a

Rigel .....	"	Rē'-jel, or Rī'-jel
Procyon .....	"	Prō'-si-on
Achernar .....	"	A-ker'-nār
Beta Centauri .....	"	Bā-ta Sen-tā'-rē
Altair .....	"	Al'-tāre
Betelgeuse .....	"	Bet'-el-gerz
Aldebaran .....	"	Al-deb'-a-ran
Alpha Crucis .....	"	Al-fa crū'-sis
Spica .....	"	Spī'-ka
Antares .....	"	An-tā'-rēz
Pollux .....	"	Pol'-uks
Fomalhaut .....	"	Fō'-mal-o
Deneb .....	"	Den'-eb
Beta Crucis .....	"	Bā-ta crū'-sis
Regulus .....	"	Reg'-u-lus

There are six stars among the twenty-one brightest which are so far south that they cannot be seen in our latitude. But they are important and beautiful, and it may be worth while to speak of them briefly before we pass on to the others of this bright company with whom we hope to have a friendly acquaintance. These six stars are Canopus, Alpha Centauri, Beta Centauri, Alpha Crucis, Beta Crucis, and Achernar. In some parts of Florida and Texas, Achernar can be seen low in the south for a short time in the evening, and Canopus, which is a little farther north, can be seen a somewhat longer time and higher up in the sky. All six of these stars can be seen in Cuba. It is to be regretted that they must remain hidden from us, for there are special reasons why some of them would interest us, aside from their great beauty.

Canopus, for instance, is far brighter than any star visible in the northern hemisphere save one, the brilliant star Sirius, and Canopus is so much farther away than Sirius that it must be many times brighter and really radiate many times more heat and light. Canopus is, in fact, one of the brightest stars in our galaxy; for all its superior brilliancy, it is so far away that light requires 230 years to reach us. Sirius, though it outshines Canopus to our view, is comparatively near since light takes only seven years to reach us.

Incidentally, it may be interesting to know that Canopus is credited with having run away with one of the Pleiades. If he did the deed, he has succeeded in concealing the proof of it, for the lost Pleiad has never been seen elsewhere since she was reputed to have disappeared from her place among her six sisters.

Alpha Centauri is distinguished as being, so far as is now known, the nearest star to us—with the exception of our sun. It is not near enough for any inconvenient neighborliness, as we shall find later on; Alpha Centauri, itself a double star, has a third, fainter component, possibly slightly nearer than Alpha.

This faint companion star is sometimes called Proxima; but the name has not come into any general use, nor does it seem very desirable that it should. For all our feeling that so important a star ought to have an individual name, Proxima does not seem the right choice. It may well be that in the course of time other stars will be found to be nearer to us than Alpha Centauri, and the name would then have no appropriateness. Alpha Centauri also bears the name Rigil Kent.


Beta Centauri, sometimes called Hadar, is a brilliant star, not quite so bright as Alpha. The two stars lie very near together. The distance between them is about four degrees. This is a little more than the distance between Castor and Pollux and not quite so much as that between the two pointers in the Dipper. Alpha Crucis, almost as bright, lies not far away, and the three stars make a brilliant spectacle in the southern sky. Achernar, but little fainter than Alpha Centauri in brightness, is the only very bright star in its section of the southern heavens.

The other sixteen of the brightest stars are all visible to us, and are the ones we shall seek to know first. After they are well identified it will be easy to point out others, less brilliant, but perhaps not less significant.

To find any of these stars only two things are necessary to be known as a basis from which to work: the constellation known as the Big Dipper, and the North Star (Polaris is its name).

In the strictest sense, the Big Dipper is not a constellation at all, but an "asterism," a group of stars that forms part of the large constellation, the Greater Bear. The Big Dipper consists of seven principal stars, all very bright, but none of them among the twenty-one brightest, grouped so as to resemble the outline of a dipper.

This star group is so conspicuous and, hence, so easy to find, that it is known to most persons. The Big Dipper is one of the few star patterns that really look like the things after which they are named. It lies in the northern heavens, and in our latitude is above the horizon all night, so that it can always be seen




THE BIG DIPPER

when the skies are clear. The two stars on the outer side of the bowl are called the "pointers," because a line drawn through them from bottom to top will point towards the North Star, and, if extended to about five times the distance between the two stars, will end, not exactly, but yet so nearly at Polaris that there will be no mistaking the star.

The North Star marks the center, or pole, around which the celestial sphere revolves, and the pointers always hold the same relative situation to it no matter how the dipper is turned with relation to us. The North Star and the Dipper as they appear to an observer on the earth in our latitude at the different seasons of the year have the relative position shown in the illustration.

Inasmuch as all the stars seen in the northern hemisphere of the heavens revolve around the point marked by the North Star, their positions relative to one another are always the same. So if one bears this in mind, it will be just as easy to find them at one time in the year, or in the evening, as at another. The positions given in the diagram of the Dipper are as the stars composing it appear early in the evening at the seasons named. They make a circuit of the pole each twenty-four hours, so that if one watches all night he can see them in at least three of these positions each night in winter and in two of them during the short nights of summer. Into the other positions they come, of course, in the daytime.



#### RELATIVE POSITION OF THE NORTH STAR AND THE DIPPER

The distance between the two pointers is about five degrees; and this is an important thing to keep in mind. These known five degrees may be used as a unit of measure in finding other stars as described in this book. If one cannot carry the distance between the pointers in his mind from one part of the skies to another and thus make comparisons of distances that will be approximately correct, he can take a stiff piece of card-board, or an adjustable ruler, and, holding it out at arm's-length, adjust it so as to exactly cover the space between the two pointers. He will then have a fairly accurate measure of five degrees in the



sky and will probably be surprised to find that it will require a little less than three inches on his ruler to cover it. This will be of great assistance in marking off approximate distances on the surface of the heavens as it appears to us, and will enable one to follow the directions given in these pages. It will be a still further convenience to make a measure from two to four times as long, thus covering ten or twenty degrees. In adjusting these measures it will be found necessary to sight with one eye only, keeping the other closed. One cannot cover a star from both eyes with a ruler at only an arm's-length away.

While these rough guides are conveniences to one entirely unaccustomed to observing the skies, a very little experience will soon enable one to estimate distances from star to star by the eye alone with sufficient accuracy to identify any conspicuous object in the heavens.

In looking for the brightest stars one may sometimes find himself confused by four bright stars that are not, after all, true stars, but the planets Venus, Jupiter, Saturn, and Mars. Stars are suns and shine by their own light, as our sun does. Planets are dark bodies and shine only by light reflected from the sun. If the planets were as far away as the stars, they would be entirely invisible, and all the differences between them and the stars which I am about to enumerate are due entirely to the comparative nearness of the planets.

There are eight planets besides the earth, but the four already mentioned are the only ones likely to be mistaken for true stars and thus prove confusing to the ordinary observer. Uranus is barely visible to the naked eye. Neptune and Pluto can be seen only with a telescope, and the other (Mercury) on account of its nearness to the sun is rarely seen at all by the ordinary observer, though when it is visible it appears as a rather bright star.

The first two, Venus and Jupiter, are at certain times brighter than the brightest stars. Venus sometimes shines so that it casts a shadow and sometimes can be seen, like the moon, in daylight. Saturn is usually as bright as a first-magnitude star, and Mars varies so that it sometimes equals Saturn in brilliancy and sometimes is not so bright. They all move about in the heavens with seeming irregularity, and when they are in the region of any bright star may easily be mistaken for it by an untrained observer. A little careful observation, however, will show that they

are not as the other stars. They are not fixed in their places, and a few nights of watching will make this apparent. They do not twinkle as the stars do, but shine with a steadier light. They are not mere points of light, as the stars are. Telescopes show them as rotund solid bodies, though small in comparison with the other two bodies of which we can see the outlines: the sun and the moon. A very little watching will make one familiar with their appearance so that he can always tell Venus and Jupiter at a glance and usually Saturn and Mars.

Their positions in the heavens are hard to describe definitely because the planets are constantly changing. They do, however, keep within a certain limited path, which is within a few degrees of the path that the moon follows as it threads its way among the stars each month, and at some time during the month it passes near each planet. This is sufficient to indicate that the planets are never found very far south or very far north.

Venus has still the further limitation of never being more than about forty-eight degrees away from the sun. When it is in the evening sky it sets never much more than three hours after the sun, and when it is seen in the morning it rises never more than about three hours earlier than the sun. The other planets within their limited path already described rise and set at various times within twelve hours of the rising and setting of the sun.

The planets have each a distinctive color, which will assist somewhat in identifying them. Venus is white; Jupiter is a little tinged with yellow, Saturn is decidedly yellow, and Mars is fiery red.

There are many interesting things to observe about the planets, which are a study in themselves. This brief outline of their appearance and locations in the sky will be sufficient to prevent one from confusing them with the stars, which are the special objects of our quest at present; and with the North Star and the Dipper well in mind, we are now ready to be introduced to the brightest of them.

### III

## Capella

CAPELLA is a charming star that has the distinction of being situated nearer to the North Star than any other of the brightest stars. To find it, draw a line forty-five degrees long from Polaris at a right angle to the line from the pointers to the pole and running in an opposite direction from the handle of the Dipper. It is also pointed out by a line starting at the star that marks the bottom of the Dipper on the handle side and running thence about half-way between the two pointers, for a distance of about fifty degrees, or to the first bright star, which is Capella. If these directions are even approximately followed Capella can be discovered, for there is no other star in its neighborhood that in any way compares with it in brightness.

Capella is one of the most brilliant stars in the heavens. Capella, Vega and Arcturus are the brightest stars in the northern hemisphere of the skies. Sirius, as we know, always shines with greater brilliancy than any other star; but Sirius lies south of the equator and does not belong among the stars of the northern hemisphere, although in our latitude it, fortunately, can be seen.

Being so far north, Capella is above the horizon more than twenty hours out of each twenty-four, so that it can be seen at some time in the night every month in the year, and one comes to be on most familiar and friendly terms with its bright face. Its first appearance in the skies of the early evening is in August, when it rises about ten o'clock during the first half of the month; and it remains a constant ornament of the nights until the latter part of the following June. July is the only month in which Capella does not show its face at some time before midnight; and even in July, if one watches closely late in the month, he can have glimpses of the star following the evening sun at no great

distance and almost lost in the sunset glow. In October it rises just as the sun is setting.

Capella rises almost exactly in the northeast and swings in a fine long circle over the heavens during its twenty-hour journey to the northwest, where it sets for a brief time, and then rises again in about four hours.

When you watch the birds congregating in noisy flocks in the morning for the fall migration, and in the afternoon look for the first fringed gentians, look for Capella in the northeastern sky in the evening. When the trees are bare and the berries are wrapped in ice and snow, so that the winter birds greedily gather in what bounty you throw to them, you will find Capella shining almost directly overhead early in the evening. During the spring months, when the air is full of the stir of the awakening earth, and other stars are demanding our attention to their return in the east after long absences, Capella is hurrying on towards the northwest, no longer charming us with its novelty, but still as bright and fair as ever and ready to fill its place in the brilliant gathering of the stars of spring.

Capella is yellow in color like our sun and has about the same surface temperature. Its spectrum is very similar to that of the sun. The similarity in temperature only means that it has about the same surface brightness. Capella, however, is a giant star and our sun is a dwarf, Capella radiating more than 100 times more light than the sun. Its distance is so great that its light requires about 46 years to reach us in its journey through space, though travelling, as light does, with a swiftness of 186,000 miles a second. At the distance of the nearest star, our beautiful big sun would shine as a star no brighter than Polaris. But Capella is ten times farther away than the nearest fixed star, and still shines as one of the brightest of the bright stars. If Capella were inhabited, our sun would appear to its people as one of the faintest stars that we can see with ease. But it is not inhabited, for it is a very hot, rarefied and expanded body, as much as forty times rarer than our sun, and like the other bright stars, no place for the existence of life as we know it.

It has been discovered in recent years that Capella is not alone as it makes its journey through the black void. Like so many other stars, it is what is known as a binary system, or a double star, and has a companion which lies so near it that they circle

around each other in a little more than a hundred days, or less than one-third of the time that it takes the earth to go around the sun.

Capella and its companion have never been seen separately even with the assistance of the strongest telescope. But the wizard spectroscope has penetrated their family secret and by its means astronomers have learned that they are two immense, fiery bodies of nearly the same size and perhaps a hundred times more voluminous than the sun, whirling with incredible speed about each other. The diameter of the orbit over which the two stars spin is so large that we could string more than a hundred suns along it and still have room left for a few more shining beads of the same sort.

Since it has been known that Capella is a double star it has been suspected through at least one telescope of being oval instead of round, showing that the two stars have been caught at a place in their orbit where they appeared side by side to our view.

These are interesting facts that science has to tell us about Capella; but the star that fixes itself in our regard is the fair, golden, bright Capella that decks the sky in its season. We follow it in its course visible to us across the heavens, we joy in its beauty and find its gentle light no less gentle on account of its being the composite of the great orbs we know it is.

Capella is at present receding from us at the rate of about twenty-three miles a second, or more than a million miles a day. At first thought this rather makes us fear that we are in danger of losing this beautiful star from view. But it will be many thousands of years before even under such speeding away there will be any appreciable change in its brightness.

The constellation of which Capella forms a part is called Auriga. It is a five-sided figure somewhat in the shape of a shield. It can be easily recognized as soon as Capella is located. It will be described among the constellations later on.

## IV

### Arcturus

**E**ARLY in the evening in March there may be seen rising a little north of east, another bright star that is among the choicest of the heavenly bodies. This is Arcturus. To find it begin at the end of the handle of the Big Dipper and, following the curve of the handle, extend a line about thirty degrees, or until you come to the first brilliantly shining star. There is no other very bright star in this region; so the line cannot point you to any other than Arcturus, the self-same star whose bright beams flowed down more than twenty-six hundred years ago upon the patient Job and is named by him along with other wonders of the divine creation.

When one has learned to know the ways of Arcturus and cares to greet him as he first comes to take his place among the evening stars in the spring, one begins about ten o'clock in the evening in February to watch the handle of the Dipper, so clearly, to the expectant observer, pointing to something important just below the horizon and then slowly rising until it has pulled into view the steady shining face of Arcturus. A less attentive observer is more likely to notice the star in March and April, when the Dipper is almost directly overhead and is turned as neatly upside down as if it were on the pantry shelf. Its handle, though, still points to Arcturus, now climbing the eastern skies composedly but rapidly, and making way for the other bright stars of the spring that will bring on the scene a stellar setting of their own.

Arcturus rises on the first of March about eight o'clock in the evening, and during this month, when the horse-chestnut buds are swelling and the elm-trees are putting forth their first brown blossoms, and the hawks and owls and crows are prospecting for nesting sites among the high trees, Arcturus, as if impelled by

the onrush of spring, is returning four minutes earlier each evening until at the beginning of April the star rises just as the sun sets.

This is really the month of its greatest glory. It shines all through the evening in the eastern heavens, bright even when the moon is full, and, fitting in with all the other aspects of nature, gives a splendid close to the splendid days of late April. What more gracious day's progress in beauty could there be than to travel with the eye from the cheerful hepaticas dotting the soft ground among the trees to the round, white, silent blossoms of the dogwood fringing the late April woods and thence, when evening falls, to the bright yet gentle light of Arcturus in the sky, announcing the end of the purple twilight.

By the middle of June, Arcturus may be found during the early evening almost overhead, and it remains in this region during July. After this, one sees plainly that Arcturus is traveling towards the northwest, sinking lower and lower each month until, near the last of November, the pointing handle of the Dipper (the Dipper being then right-side up and below the pole) shows us that it has gone, almost at the same moment with the sun, below the northwestern horizon. It is in the evening skies no more until the next spring. If one wants to see Arcturus in December or January, he must look for it in the east between midnight and dawn. It shines for us at some time between sunset and midnight almost ten months in the year. It requires about fourteen and a half hours to complete its journey from rising to setting each day.

Arcturus and Capella are so nearly equal in brightness that astronomers have differed as to which outranks the other. To my own eye Arcturus outshines Capella, and on asking various inexperienced persons for off-hand opinions as to the relative brightness of the two stars, I have invariably had an answer in favor of Arcturus. The best authorities now agree with this impression.

Compare Arcturus with Capella and you will see that the colors do not precisely match. Arcturus, as one can plainly see, is a little more tinged with red. This means that it is cooler and the surface not quite so bright as that of Capella or the sun.

These red giant stars are in a very interesting stage of their astronomical development. They are stars that, having exhausted

the primary stellar fuel, hydrogen, are now relying upon secondary sources in an effort to escape extinction. Their centers are much hotter than the sun and this extra heat has caused the stars to expand until they are bloated in comparison with the sun.

The total brightness of Arcturus is almost a hundred times that of our sun. Hence, since the source of fuel is very limited, this star and other similar red giants are expending their energy at an enormous rate, which they cannot keep up for long. After a few thousand years these stars may explode, the remaining fragments condensing to form white dwarfs, like the companion of Sirius.

Light takes about 37 years to reach us from Arcturus.

These facts are, perhaps, not what we most want to have in mind when we gaze into the face of Arcturus with its clear, steady light, but they must have their part in the sum of its general characteristics and some influence in determining our feeling towards the star.

Arcturus has the distinction of having the most rapid motion of any of the brightest stars and it is among the swiftest moving of all the stars visible to the naked eye. Enormous body as it is, we know that it is bowling along through space at the rate of between two and three hundred miles a second. It has been doing this for untold centuries, and yet it does not change its position to our view a distance of more than one-tenth of the apparent diameter of the moon in a century. This is, however, an enormous change for a "fixed" star and will, in the course of ages, make an important difference in the position of Arcturus among the other stars.

It is pleasant to know that this bright star is approaching us, though it decreases the distance between itself and us only at the rate of two and one-half miles a second, a very loitering gait in comparison with the speed of stars in general. The most that it can mean to us is that Arcturus will not grow less bright in our skies because of increasing distance; for at this rate of approach untold centuries must pass before the distance between us and the star is lessened so as to make any appreciable increase in its brightness. Such increase, indeed, possibly never will be made. For, as previously noted, Arcturus is growing old and



may well alter its appearance before changing distances have much influence on its apparent brightness.

Arcturus lies in the constellation Boötes and is, of course, its brightest star. The name signifies "Watcher of the Bear," a title derived from its proximity to Ursa Major.

## V

### Spica

**A**NOTHER bright star, and one that comes into view not long after Arcturus, is Spica. It is not so bright as Arcturus, but is still a conspicuously brilliant star. Among the twenty-one brightest it is the fifteenth in order of brilliancy. To find Spica easily extend the line from the end of the handle of the Dipper through Arcturus and about thirty degrees beyond, still following the curve. It is about as far from Arcturus as Arcturus is from the handle of the Dipper. A straight line beginning at the pointer nearest the pole, running through the star at the bottom of the Dipper on the handle side, and extended about sixty-five degrees farther on, will also point out Spica.

Spica rises a very little south of the exact eastern point on the horizon and becomes visible about an hour later than Arcturus. It is the only very bright star in the southeastern part of the heavens during the early spring months, and one cannot easily fail to identify it even without the specific directions for finding it that are here given. It first attracts one's attention in the early evening about the middle of March, at which time it rises about eight o'clock and by nine has cleared its face of the haziness of the lower atmosphere and is shining with a clear white light.

Spica does not make so long a sweep across the heavens as Capella and Arcturus do. It is one of the stars that lie a little south of the celestial equator, and at the highest point in its journey across the sky Spica is still to the south of us and almost one-half of the way up from the horizon to the point exactly overhead, which we call the zenith. It takes five hours and a half to reach its highest point and as many more to travel to the southwestern horizon, where it sets eleven hours after it has risen.

Spica rises just after the sun sets in the early part of April; and during April and May it is at its best in the eastern evening sky. Its position, however, always comparatively low in the sky to our view, makes it easy to observe at all times when it is visible. During the short evenings in June Spica lies towards the southwest, where in August it sets about ten o'clock, in September at about eight o'clock, and in October about the same time that the sun goes down. It can be seen between sunset and midnight in some part of the sky from the middle of February to the early part of October. In November, December, and January it rises between midnight and dawn, so that October is the only month during which it is most of the time invisible. This is because it lies in the constellation Virgo, or the Virgin, through which the sun is passing in October, and the star is lost in the rays of the sun. Spica lies almost on the ecliptic, which is the line that marks the annual path of the sun through the heavens.

Spica's color (blue white) shows that its surface temperature is higher and surface brightness greater than that of either Capella or Arcturus. It is, therefore, a much smaller object. It is about six hundred times brighter than our sun and so far away that it takes light about two hundred and sixty years to reach us. Spica is receding from the earth at a snail's pace, compared with most stars—about one mile per second.

The name Spica means an ear of wheat, and the star marks the spot in the constellation Virgo where the Virgin holds in her hand a spear of wheat.

## VI

### Vega

**A**BOUT three hours after Arcturus has risen there will come peeping over the northeastern horizon a brilliant, bluish star which twinkles so gayly and commands such instant admiration that its entrance into view has almost a dramatic effect. This is Vega, the third of the trio of bright stars that give a May-dance around the pole. Her companions, bright Capella and steady-shining Arcturus, are already gliding smoothly over the shining surface of the sky when she, clearing the tree-tops almost with a bound, lightly springs into view and the vernal festival is on.

The point on the horizon at which Vega makes her appearance is about fifty degrees from the pole. She lacks about seven degrees of being as far north as Capella, and is nearly twenty degrees farther north than Arcturus.

To find Vega, draw a line from the star at the junction of the bowl and the handle of the Dipper to Polaris. Then draw a line at right-angles to this nearly forty degrees in length and on the same side of the Dipper from which the handle projects, and it will point out the brilliant, beautiful Vega. The star is the only conspicuous one in the region described, and no other star can be mistaken for it.


Vega is on the opposite side of the pole from Capella, but the two stars are usually seen in the sky at the same time. In the spring, when Vega decks the eastern heavens, Capella glows in the northwest. In the autumn, when Capella comes sparkling over the dark purple sky in the northeast, Vega may be seen flying with its attendant stars down towards the northwest. Each one is hidden below the horizon for the few hours that the other holds the center of the celestial stage and is at its highest point in its nightly course across the jewelled dome of the sky.

From Vega as the starting-point, you will find that Arcturus

is about one-third of the way between Vega and Capella, though a little farther away from the pole. The part that Arcturus plays in the gay spring festivities is very conspicuous, and at that time the three stars are closely associated in their splendid, twinkling circuit around the pole which it may please our fancy to watch during the first soft nights of spring. In the autumn, when Vega and Capella have changed places in the evening skies, Arcturus is below the horizon and cannot be seen.

If one is in any doubt about the identity of Vega, it can always be recognized at a glance by the fact that four dainty stars arranged in an oblique parallelogram follow it.


The position shown in the drawing is the one they always hold relative to Vega. When we see her in the east as she bright-



VEGA (IN LYRA) AS SHE APPEARS IN THE EAST

ens the evening view in the spring and early summer the four stars are to the east of, or below, her; in the late autumn and winter, when the constellation is in the western sky, the position of the stars, as it appears to us, is partly reversed, and, with Vega leading them towards the northwestern horizon, they appear as shown in the illustration.

Vega, like Capella, can be seen in our latitude at some time during the night every night in the year. One can see it even between sunset and midnight every night in the year. But to do this it will be necessary to keep a sharp lookout low in the northwest very early in the evenings in February and to turn just as sharp eyes towards the northeast between eleven and twelve



VEGA AND HER ATTENDANT STARS IN THE NORTHWEST

o'clock in the early part of March. As April progresses the star more and more asserts its supremacy in the northeastern heavens. During the latter part of this month and all of May it is singularly beautiful as it passes almost airily between us and the rich dark background so full of color characteristic of the spring-time sky.

Vega is the chief star in the constellation called Lyra, the Lyre—once fancied to be the magic lyre which in Orpheus' hands gave forth music that tamed wild beasts and chained the rivers in their courses. It is easy to get some sense of the fancy that gave the constellation its name as we watch it during the lovely spring evenings floating lightly in the sky, the parallel lines connecting its principal stars vaguely suggestive, to the willing mind, of some quaint stringed instrument that under a magic touch might send out heavenly music through the resonant air.

Early in May the star rises at about the same hour that the sun sets, and all summer long it is the gayest and perhaps the most instantly attractive star in the evening skies. It is the star so often noticed and commented on as shining with great brilliancy directly overhead between nine and ten o'clock during the hot summer evenings of July and August. When autumn comes Vega has passed the highest point in its journey across the heavens and may be seen travelling towards the northwest. Here it shines splendidly during the early evening, setting in December somewhere near eleven o'clock and in January about

nine. It is above the horizon eighteen hours out of each twenty-four and rises six hours after it has set.

Vega is one of the three most brilliant stars in the skies of the northern hemisphere; Capella and Arcturus closely match its brightness. Only three stars in the entire heavens, north or south, are appreciably brighter. These are Sirius, Canopus, and Alpha Centauri, all stars of the southern hemisphere. Sirius, as we know, can be seen in our latitude, but it lies south of the celestial equator, and is, therefore, called a southern star. The other two lie so far south that they do not ordinarily come within our view, although Canopus barely creeps above the south horizon for southern locations.

Vega is one of the very bright suns of the universe and is 50 times brighter than our sun. Its distance is such that it requires about twenty-seven years for light to travel from it to us; thus Vega is somewhat nearer to us than Capella and Arcturus. The surface temperature of Vega is less than that of Spica but considerably greater than that of Capella or Arcturus.

One of the most interesting things about Vega is that it seems to mark, or nearly mark, the point in space toward which our sun with all his planets is travelling. Vega, of course, has nothing whatever to do with this motion. It is coming nearer all the time, too. Ten miles a second is about the rate at which it is approaching us; but, as we have already seen, it will take many centuries for even such almost incredible speed as that to make any appreciable difference to us in the brightness of a star.

On some old maps Lyra appears as a Vulture. Vega is said to mean "the swooping vulture."

## VII

### Deneb

A LITTLE more than an hour after Vega has risen, a bright twinkling star will appear on the northeastern horizon at a point about forty-five degrees from the Pole Star. This is Deneb, nineteenth in the list of the twenty-one brightest stars, but a very interesting and beautiful star. It lies in one of the densest parts of the Milky Way and is conspicuously brilliant in contrast with the hazy light of the myriads of small stars that go to make up this mysterious belt of shining bodies.

Deneb is about six degrees farther north than Vega and only about one degree farther south than Capella, which, as we have seen, is the northernmost of all the bright stars. To the ordinary view Deneb and Capella seem to rise at about the same spot on the horizon, and they describe very much the same long, oblique curve across the heavens as they pass from rising to setting. Capella being a little farther north, is above the horizon perhaps half an hour longer than Deneb; but they each take approximately ten hours to come to the highest point in their path across the sky, and, consequently, in a favorable atmosphere, can be seen for about twenty hours each day, except when they are obscured by the sunlight. The highest point a star ever reaches in its daily course over the heavens is marked by what astronomers call its meridian, and when it is at this point it is said to be "on the meridian."


A line drawn from Polaris to Deneb and thence on to Vega forms a right-angled triangle of which Deneb lies in the right angle.

Deneb forms a part of a figure called the Northern Cross, which extends over the Milky Way something more than twenty degrees. The figure serves to identify the star, but it is a rather ill-formed cross and not much more like a true one than the



far-famed Southern Cross, which is so impressively described by travellers in southern latitudes and is so disappointing, on first sight, to the observer who expects accurate rather than suggestive outlines in the constellations.


Deneb is the star at the head of the upright of the cross. In the spring, when it rises in the northeast, the cross is almost upside down, and Deneb comes up last. When the cross is setting



TRIANGLE FORMED BY POLARIS, DENEK, AND VEGA

in the northwest, its position is partly reversed, so that it seems more nearly upright, and Deneb disappears last.

While this figure is commonly spoken of as the Northern Cross, it is really only a part of a larger constellation known as



#### THE NORTHERN CROSS, PART OF CYGNUS, THE SWAN

**Cygnus**, or the Swan, and this is the name used by astronomers in referring to the constellation in which Deneb is found. It was named by the ancient astronomers, and was thought by them to suggest the figure of a flying swan, the out-stretched wings being represented by the stars which form the arms of the cross. The head is marked by the star at the base of the cross, and in the tail is Deneb. The name "Deneb" means "the tail," and the star gets its name from its position in the constellation. Deneb

has sometimes been called *Aried*, an Arabic word which means "the hindmost," though the name is rarely used.

The swan represented by this constellation was supposed to be the one whose form Jupiter assumed when he visited Leda incognito; and having served the purpose of the god, what more natural than that its empty but sacred form should be placed forever in the heavens?

Although eighteen other stars (fourteen in our latitude) seem to us brighter, Deneb exceeds most of them in size and certainly in intrinsic brilliancy. Actually, Deneb is about sixty thousand times brighter than our sun. Also it is very distant, so far away that light takes about 1400 years to reach us. This means that Deneb is so many millions of millions of miles away that if its distance were expressed in figures they would be more than the mind can grasp.

Deneb is white in color, with a surface temperature similar to that of Vega. However, it is indeed a much larger star than Vega, of a class sometimes called "supergiant."

Deneb, like Capella and Vega, can be seen at some time between sunset and midnight every night in the year. It is closely associated with Vega as an ornament of the sky, because they adorn the same portion of the heavens at the same season. Following almost in the wake of so dazzling a star as Vega, Deneb may at first excite less interest and admiration than is its due; but a closer acquaintance with its sparkling face will show that its beauty is no more eclipsed by Vega than the low nodding anemones are eclipsed by the splendor of the magnolias. They each have their own individual charm.

Deneb arrives at the spring gathering in the east an hour later than Vega, but, being farther north, it pursues a longer course, and, coming leisurely to the western sky, it lingers in our view each day for three hours after Vega has gone. Deneb is particularly attractive in this part of its path during the early evening in January and February. It is then rather low in the northwest, and with Vega gone and no other bright star very near, it has a more commanding charm. In the heavier atmosphere, which induces a more rapid twinkling, the star seems to take on an accession of gayety and goes fairly dancing down behind the horizon, where it finishes its circle and appears again in the northeast about four hours later.

Deneb rises just after the sun sets about the middle of March and sets about an hour after sun-down during the early part of March.

Inasmuch as Deneb is so far distant, it is interesting to know that in the same constellation (Cygnus) and not many degrees east of Deneb, or below it as it rises, lies a star famous as being the first whose distance from us was measured. This star, which bears the unprepossessing name of 61 Cygni, is about eleven light years away. Currently (1962) it stands eleventh in the list of nearer stars, not counting the sun. 61 Cygni, which is its catalog designation, is not much more than visible to the naked eye, though a keen eye can see it on a clear night. Its position is marked on the diagram of the cross on page 32.

The star at the foot of the cross, or in the beak of the swan as one chooses to figure the constellation, is Albireo, a fine double star. It is easily resolved into two stars by a small telescope and is notable on account of the strong contrast in color between the two stars composing it. One is a deep blue, and the other a bright orange.

## VIII

### Altair

A LINE about eighty degrees in length drawn from the pole towards the south and running about half-way between Deneb and Vega will point to the brilliant white star Altair. It is almost twenty degrees directly south from Albireo, mentioned in the last chapter as marking the base of the Northern Cross. It lies just at the edge of the Milky Way; and, for any one who has learned the position of Deneb, it will be easy to find Altair by running the eye down the Milky Way from Deneb along the upright of the Cross to Albireo at the foot of it, and thence on twenty degrees, or to the first very bright star, which will be no other than Altair.


The constellation in which Altair is situated is known as Aquila, or the eagle; and Altair is the middle one of three fairly bright stars placed with reference to each other in a straight line. Indeed the name Altair means "the Flying Eagle."

The lines between Altair and Deneb and Vega form an acute-angled triangle with the short side towards the north and Altair at the apex at the south.

Altair rises about eight degrees north of the exact eastern point on the horizon and about ten degrees south of the spot where Arcturus first appears. It is the only very bright star in this region of the sky, and there will be no trouble in identifying it. One of the interesting figures that serve to mark the locality is a small diamond-shaped group of four stars lying about ten degrees northeast of Altair.


This group is the constellation Delphinus, or the Dolphin, more popularly known as "Job's Coffin." Why the figure should suggest a coffin rather than a dolphin, it is not easy to see. It hardly resembles either object.

Among the stars of the northern hemisphere, Altair is the first



ALTAIR AND ITS TWO COMPANIONS IN AQUILA


bright star to claim our attention in the east after Vega and Deneb have made their brilliant close to the gathering of the stars of spring. Capella is within two hours of setting in the north-west, Arcturus is nearing the highest point in his course across the heavens, Spica has reached the meridian, and Vega is about half-way up to the zenith, when, on the first of June, between half-past eight and nine o'clock in the evening, Altair makes its gracious entrance into the starry company. With one exception, it is the most attractive of the stars that rise during the early evening in summer. By the middle of the first summer month there is a lull in the noisy rush of the on-coming season. The dazzling blossoms of the fruit trees are scattered and almost forgotten, while the tiny green outlines of the fruit are quietly swelling towards their perfect form. The leaves on the forest trees have assumed their proper shape and are gradually expanding to full size. The old-fashioned annual roses are in their



DENEb, ALTAIR, AND VEGA

fullest glory. The blue jays have become stealthy and quiet; and most of the birds, while still in full song, have given over the frolicsome hilarity of the spring for the serious work of the nesting-time.

Then there comes a soft June evening, with its lovely twilight that begins with the last song of the wood-thrush and ends with



### THE DOLPHIN (DELPHINUS), OR JOB'S COFFIN

the first strenuous admonitions of the whippoorwill; and, almost as if it were an impulse of nature, one walks to the eastern end of the porch and looks for Altair.

It is sure to be there smiling at one just over the tree-tops, with a bright companion on either side, the three gently advancing in a straight line as if they were walking the Milky Way hand in hand and three abreast.

From this time on until the first half of December is past, Altair is in view early in the evening. Always an attractive star, it is never more charming than in September and October. It then lies due south of us and at its highest point, which is about two-thirds of the way up from the horizon to the zenith. This favorable position, added to the peculiar clearness of the autumn sky, brings out all the soft beauty of the star.

One seldom sees Altair during the first four months of the year. Along in the very first days of January a prompt lookout and a clear sky will yield a glimpse of it between sunset and seven o'clock; but it is soon lost in the glow of the rapidly lengthening days. In February, March, and April the very early



riser and the belated watcher after midnight may see the star rise, and in May, as the month progresses, it becomes a more and more familiar figure in the east.

There is nothing spectacular in the advent of Altair, and one does not watch for it with the same eagerness that he feels when Arcturus and Vega are nearly due. It comes gently into sight, and almost before one realizes it the star has become a part of the evening view. Arcturus and Vega suggest the bustling coming of spring. Altair reminds us of the quiet approach of summer.

It requires about thirteen hours for Altair to pass from its rising to its setting; six hours and a half to reach the meridian, and six and a half more to finish its diurnal course. Remaining as low in the sky as it does, one can see it and yet remain in a comfortable posture during its entire passage across the heavens. This adds greatly to its charm; for any star is, in itself, most beautiful when in the highest and clearest part of the sky. Only the difficulty of seeing it there may detract from our pleasure in it. One cannot come into any great fondness for a star that he sees only by snatches. The stars that we love best are the ones into whose faces we can look for an hour at a time, if our fancy so leads us. It is truly star-gazing that brings us into any great delight in a star.

Altair is white or yellowish white. Its surface temperature is somewhat less than that of Vega but still higher than that of Capella. Its true luminosity is about eleven times that of the sun. Light takes about seventeen years to reach us. In this respect, Altair is not one of the far distant suns, but is comparatively near us. The star itself is approaching us at the rate of about sixteen miles a second or more than 200 millions of miles a year. Yet even with this marvelous speeding toward us, and in spite of its comparative nearness, considerably more than 100 thousand years would be required for Altair to reach the place in the universe that we now occupy.

## IX


### Antares

**A**BOUT ten o'clock in the evening of the longest day in June is, perhaps, the most favorable time to look for Antares—a beautiful star in a beautiful constellation. It then lies due south and at the highest point in its diurnal path across the heavens, which never takes it higher than a little more than one-fourth of the way from the horizon to the zenith. A line about one hundred and sixteen degrees long drawn directly south from Polaris and passing just half-way between Arcturus and Vega will point it out, lying in the heart of the constellation Scorpius, which is flung out over the southern sky in a fine sweep that wins admiration from the most casual observer.

Antares is also indicated by a line drawn at right angles to the line connecting Altair and its two companions and extended about sixty degrees, or to the first brilliant red star, Antares, the only one of its kind in this most interesting region of the skies. An almost equal-sided triangle about fifty degrees long on each side is formed by the lines connecting Antares, Vega, and Arcturus, which is a still further guide to the location of the star.

Antares is one of the southern stars that lie south of the celestial equator, no matter whether they are visible to us or not. It is, with one exception, farther south than any other of the very bright stars (stars of the first magnitude) that can be seen in our latitude. It rises in the southeast and has so short a course over which to pass, in our view, that it reaches the half-way point, or the meridian, in about four hours and a quarter.

Antares comes into view two hours and a half later than Vega, an hour and a half later than Deneb, and nearly an hour earlier than Altair. But it is so much farther south than Altair that it seems more aloof from us and it scarcely emerges from the hazy



lower atmosphere until Altair is shining gayly and has already become one of the important figures in the eastern summer sky.

While the latter part of June and all of July are perhaps the best times to see Antares at its brightest, it is an interesting star to observe much earlier in the season. If one has an unobstructed view of the southeastern sky down almost to the horizon, he will

note the appearance of Antares as early as the middle of May between nine and ten o'clock in the evening. Before one has really seen the star, he becomes conscious of a ruddy glow low in the southeast that at once fastens the attention. It is the face of Antares, whose red light shining through the heavy atmosphere is so diffused that it gives a rosy effect to the sky for a considerable distance around the star, like a miniature presentment of the sun as it rises on a hazy morning. This is all the more noticeable because it is so in contrast with the usual colors of early May. They are all of the more delicate and colder sort, from the pinkish-lavender and white of the young leaves that shimmer over the landscape to the pinkish-purple of the sky and the dazzling white and blue of Deneb and Vega and a few other stars that belong particularly to this season.

The fact that Antares, when it is visible at all, is always so low in the sky and so clearly before our eyes when they are turned towards the south, makes it a constantly conspicuous figure in the heavens during its entire stay. It is fortunate that this is so, for the stay of Antares is short. It is visible in the evening between sunset and midnight for only six months, and during only one of these months (June) is it visible all night. In November and December and most of January it is above the horizon only in the daytime, and in February, March, and most of April it is not high enough to be clearly visible until after midnight. In the early part of June it rises just as the sun sets.

I am not sure that the color of Antares is any reason why it should be associated with the blooming of red flowers; but I find in my journal of out-door things that I have unconsciously made the association. One entry, on June 30th, says: "Antares is shining splendidly to-night and rivals in color the wild red lilies that were blooming to-day in the coppice down beyond the spring." Another entry, on August 5th, says: "We found the first cardinal-flowers to-day," and a little farther on, "Antares is almost as red as a cardinal-flower to-night." It is certain that the wood lilies bloom when the star is at the culmination of its summer glory, and the cardinal-flower sends up its flaming stalks when the star is speeding after the sun and sets about midnight. Its bright red color, too, is accentuated when Antares is near the setting-point in the southwest, just as when it was rising in the southeast.

Although Antares is almost strictly a summer star, it is a very significant object in the southwestern heavens during September and October and is often commented on at that time by persons who have not noticed it earlier in the season. The dark background of the autumn sky throws the flaming face of the star into relief, and its redness is still further intensified by the white sweep of the Milky Way, which is then very bright and which lies within a few degrees of Antares.

The constellation of which Antares forms a part (Scorpius) represents the scorpion that stung Orion, and it is the "hissing snake" from which Spenser represents Orion as "flying fast." The two constellations are so placed in the skies that Orion dips below the western horizon just as Scorpius rises in the east, and the mighty hunter does not return until the Scorpion is gone. They can never be seen at the same time.

The name Antares signifies "Rival of Mars." And indeed the red color of these two objects is most striking. It is particularly so when, every couple of years, Mars in its course through the ecliptic chances to be in Scorpius. For then the star and the planet stand out in splendid contrast.

Antares belongs to that class of stars known as the red giants. It is even more distended and swollen than Arcturus. Its redness indicates that the star is cooler, with a less luminous surface than Arcturus has. Actually it is about five thousand times brighter than our sun and very much larger. If Antares were set in the position of our sun, the earth would still be somewhat below the surface. Light takes about 400 years to reach us from this star.

Red giant stars, like Antares, are believed to be objects in their old age. Most stars derive their energy from the fusion of hydrogen atoms into heavier substances like helium. But when a star has exhausted most of its hydrogen, the internal core tends to contract, grow hotter, so that the star can for a very short time live on other energy sources. The outer layers swell up and the star becomes very luminous. But after some thousands of years of this spendthrift activity, these sources, too, become exhausted. The star has nothing left to do but collapse on the inside and explode on the outside, becoming a nova, or new star. Gravitation will then collect the fragments into a small,

feebly shining star of the type known as a white dwarf, like the companion of Sirius.

Antares is one of a binary system. Its companion is remarkable for its greenish color, which forms a strong contrast with the scarlet light of the larger star. The companion can be seen, however, only with a fairly large telescope, partly because of its proximity to the primary and also because it is three magnitudes fainter.


## X

### Fomalhaut

STILL lower in the southern sky than Antares and in a comparatively starless region shines Fomalhaut, the lonely, the eighteenth star in point of brightness among all the stars in the heavens and the thirteenth among those that can be seen in our latitude. It is so completely isolated from other bright stars that it needs no special line to point it out. But if the line that leads from the pointers to the Pole Star were extended beyond the pole and on directly south about one hundred and twenty degrees, it would end near Fomalhaut. This will be an easy line to trace during the months when Fomalhaut is visible, for the pointers will then be shifting from a position west of the pole to one directly under it, and the line will have a free path across the heavens. The star may be easily recognized without any other description than that it is at the end of a zigzag line of small twinkling stars which runs across the southern sky in the autumn and points unmistakably to Fomalhaut. The small stars represent the stream of water that Aquarius is pouring from his urn. The stream runs to the constellation known as the Southern Fish (*Piscis Austrinus*), and Fomalhaut means the mouth of the fish.

There is a calm dignity about this star that lends much interest to it. It comes gently into view far down in the southeast in August, but with so little of a flourish that one scarcely notes its presence until along in early September. Then, when the days are growing shorter, some evening, just after dark, one sees it, a conspicuous, impressive object serenely trailing along over the small arc of its circle in the south with no companion near it, and, apparently, no need of one to add to its splendor.

On early acquaintance the loneliness of the star, added to the sombre signs of approaching autumn, sometimes gives one a touch of melancholy. But its aspect when more familiar soon



## FOMALHAUT

comes to suggest only sweetness and serenity, and a lover of Fomalhaut feels that a sustaining light has gone when, during the last of December, this beautiful star sinks gently down in the southwest and disappears from the evening sky not to return for more than seven months. One loves to recall that Fomalhaut makes a lonely but majestic sweep high in the heavens of the southern hemisphere: a sweep somewhat larger than Arcturus makes in our skies.

Fomalhaut and Capella rise at almost exactly the same minute; but the path of Fomalhaut across the sky is so short that it has finished its course and vanished more than two hours and a half before Capella has reached its half-way point, the meridian. Rising together as the two stars do, one seeks to continue some close association between them; but they have nothing more in common. In the course each has to run and the joy each has to give they are as widely divergent as two royal babes born at the same moment on different sides of the globe.

Fomalhaut is one of the four "royal" stars of astrology. The others are Regulus, Antares, and Aldebaran.

Only eight hours are required for Fomalhaut to pass from rising to setting, and at its highest it is never quite one-fourth of the way up from horizon to zenith. It is the farthest south of any very bright star visible in our latitude.

Fomalhaut is distinctly a star of the autumn. When it first



appears in August the nights have already begun to grow longer, the birds have ceased to sing and are seeking the deeper shadows of the woods in which to shed their summer suits, and the quiet of autumn is everywhere stealthily creeping upon us. At the time of the equinox, in September, the star rises about an hour after night falls and is on the meridian between ten and eleven o'clock in the evening. About the middle of October Fomalhaut rises as the sun sets and shines until two o'clock during the first white frosty nights. In November and December, when even the most hardy insects have stopped their shrill notes and the winter stillness has fallen upon us, a glance towards the southwest will discover Fomalhaut, still placid and alone, on its way to adorn the warmer southern skies, its serenity in no wise disturbed and its beauty in no wise eclipsed by the more spectacular entrance of the brilliant winter stars now gathering in the east.

Notwithstanding that to our view Fomalhaut is surpassed in brightness by seventeen other stars, it actually gives out about fourteen times as much light as the sun. While not one of our nearest neighbors among the stars, it is still not so very far away, for its light reaches us after a journey through space of only 22 years. There are only four other stars among the twenty-one brightest that are known to be nearer to us.

Fomalhaut has no companion star. It is not only in a more or less starless region to our view, but its loneliness seems to be actual.

I have preferred the pronunciation Fo'-mal-o as being more melodious than Fo-mal-hoot', a pronunciation sometimes recommended by the authorities.

.

## XI

### Aldebaran


THE stars thus far described have a particular connection with the spring, summer, and autumn months. Capella, it is true, shines throughout the winter; but it can be seen while the evenings are pleasant, and as it is in the most convenient position for observation in the spring and fall, it belongs, in large part, to the out-door season. There are, however, six brilliant stars that are distinctly of the winter. They are not only of the same season, but in a way they belong together, for they are in constellations that join each other and spread out in a splendid panorama across the southern skies, making the most dazzling display of fixed stars that any part of the heavens affords. These six stars are Aldebaran, Betelgeuse, Rigel, Pollux, Procyon, and Sirius.

Of these six stars Aldebaran is the first to appear. Along in September, a very little north of east it shows its fiery face above the horizon with such unmistakable individuality that it catches the eye of even the least observing. It appears at a point about thirty degrees farther south than Capella and rises about two hours and a half later than that bright star. But it is hardly less conspicuous, for though its actual brightness is rated nearly a magnitude lower than Capella's, it glows with a rosy light that demands recognition and at once pronounces it one of the most important heavenly bodies.

Aldebaran lies at the open end of a bright V-shaped cluster of stars known as the Hyades.

The cluster is in the head of the great constellation Taurus, or the Bull, and the star itself marks the fiery eye of the enraged bull as he lowers his head for a lunge at Orion.

It is difficult to miss finding Aldebaran if one watches in the region indicated, during late September or early October, for the first bright star that rises after Capella. The locality is still



THE HYADES

more distinctly marked if one chances to know the shimmering light of the Pleiades, which are in the same constellation as Aldebaran and constitute the one group of stars that is usually familiar to the youth who celebrate Hallowe'en. Aldebaran shoots its ruddy face above the horizon just an hour after the hazy little dipper of the Pleiades has appeared and the star is then to the east of, and, hence, almost directly under, the Pleiades. This following of the Pleiades is what gives Aldebaran its name, which means "the hindmost," or "the follower."

Aldebaran shines in the evening skies for about eight months in the year. It becomes a fixed ornament of the eastern sky about the last of September, when it rises between nine and ten o'clock. The ruby Antares has been below the western horizon for nearly an hour when the still brighter ruby Aldebaran comes into view in the east. The two stars can never be seen at the same time and they do not at all resemble each other except in color. Even in this respect they differ, for Antares is a deeper red than Aldebaran, though not so bright a star. But as one watches the slow sinking of red Antares in the southwest, and gives a sigh of regret when he has entirely disappeared, it seems but a natural association that draws one's eye towards the east to watch with eager-

ness for ruddy Aldebaran to give that touch of color which the sky seems to lack.

In astrology these two stars were known as "royal" stars, and there is a whimsical suggestion of royalty in them now as the one slips away from our admiring gaze and we turn to hail the coming of the other. "The king is dead! Long live the king!"

In his section of the sky Aldebaran reigns throughout all the lovely autumn evenings, with beautiful Capella in her own realm to the north of him and Fomalhaut far to the south. His near associates are in the bright groups that I have called the winter stars, and they appear in quick succession after him.

From the time Aldebaran first appears until late in December it may be seen in the eastern heavens in the early evening. It rises just as the sun sets near the first of December and reaches the meridian seven hours later. In January and February, Aldebaran is at its highest point early in the evening and is then a little more than two-thirds of the way up to the zenith. At the same hours in April it is conspicuous in the western sky, and by the end of May it has disappeared. In June it is above the horizon only in the daytime, and in July it can be seen not long before the sun rises.

Aldebaran is the thirteenth star in order of brightness in the entire heavens and the ninth among those seen in our latitude. It is what is known as a standard first-magnitude star. It gives to us about one 90-billionth as much light as the sun, but at the same distance from it as we are from the sun we would get from it about 100 times as much light as we do from the sun. It requires approximately 70 years for the light of Aldebaran to reach the earth, which means that it is nearly 400 trillions of miles away. It is increasing this distance at the rate of about 30 miles a second; but even at this rate it will require more than 10,000 years to add another trillion of miles to its distance.

Aldebaran lies in the path that the moon follows as it crosses the heavens, and is frequently obscured from our view by the moon's passing between it and the earth. Such an intrusion of the moon is called an occultation, and is an interesting phenomenon when the subject occulted is either a bright star or a planet. A star, in such a case, disappears behind the moon instantly, because a star shows no sensible disk. A planet disappears gradually.

Aldebaran has an extremely faint companion star, which can be seen only with a powerful telescope. Aldebaran also belongs to the class of old, dying stars—a red giant like Antares.

## XII


### Orion's Bright Stars

**N**EXT to the Great Dipper, Orion is probably the most widely known of all the constellations. It contains more bright stars than any other single group, and they exact all the more immediate admiration because they are clustered so closely and symmetrically as to form a set figure of dazzling jewels—a veritable “sunburst” of “diamonds in the sky.”

To the ancients this figure, as its name indicates, represented the giant Orion placed in the heavens in heroic attitude facing Taurus, the mighty bull, who, with fire darting from his eye (marked by Aldebaran), is about to make an onset with his splendid long horns. Orion stands with his right arm uplifted and his great club in the air ready to strike. Over his left arm hangs a lion's skin, which he holds up as a sort of shield before him. With a little play of fancy it is not difficult to see in this constellation a colossal figure, splendidly poised and full of action. The accompanying diagram shows the figure as outlined by the stars.

Orion not only includes an unusual number of bright stars, but it has the further distinction of being the only constellation visible in our latitude that contains as many as two stars of the first magnitude, or highest degree of brilliancy. These two stars are Betelgeuse and Rigel. Betelgeuse marks the right shoulder of Orion, and Rigel is in the left foot. In the strictest sense, Betelgeuse signifies armpit and Rigel the left leg.

The first of these to appear is Betelgeuse. It pushes its red face up over the horizon about an hour and a half after Aldebaran has risen, and is only about nine degrees farther south than Aldebaran. It is of about the same degree of brightness as Altair, but it has not the same cheerfulness of aspect. On the contrary, there is a decided suggestion of sombreness in its dull and comparatively untwinkling face.



ORION

The color of Betelgeuse is deep red. It is, therefore, a conspicuous member of the group of red giants—like Antares and nearby Aldebaran, stars nearing extinction. This period of stellar evolution is probably responsible for the fact that the star fluctu-

ates in brightness, for Betelgeuse is one of the class known as irregular variable stars. In other words, it passes irregularly from its brightest to its dullest stage of lucidity. Occasionally it attains the brilliancy of Vega or Capella; sometimes it is no brighter than Spica or Antares. But it always remains bright enough to be a first-magnitude star.

About fifteen or twenty minutes after Betelgeuse has appeared Rigel comes twinkling into view. It is the brightest star in Orion and lies about sixteen degrees south of Betelgeuse. The celestial equator runs just about half-way between the two stars; so that Betelgeuse belongs to the northern hemisphere and Rigel to the southern. With Aldebaran, these two stars form a triangle, the sides being from sixteen to twenty degrees in length. Rigel is exactly south of Capella; hence, a line drawn from Polaris through Capella and extended a little more than fifty degrees will end at Rigel.

In point of brilliancy Rigel ranks next to Capella. It is a very bright blue-white star, as dazzling as a small electric light and as sparkling as a fine diamond. But with all its fair beauty Rigel does not seem as much of a personage in the assemblage of stars as sombre old red-faced Betelgeuse; and in looking at them I am always conscious of feeling a somewhat closer friendliness for Betelgeuse than for Rigel. This, however, is purely a personal fancy.

With the exception of Deneb, Rigel and Betelgeuse are the most distant of all twenty-one stars of the first magnitude. From Rigel light takes 1300 years to reach us. From Betelgeuse it takes 500 years.

From the fact that Rigel and Betelgeuse, despite their distance, are still first-magnitude stars, we must conclude that they are intrinsically extremely bright. Enormous Betelgeuse, red and cool, is about 17,000 times brighter than the sun. Rigel, blue-white and extremely hot, though very much smaller than its nearby red sister, is 150,000 times more brilliant than the sun.

Although Rigel rises about fifteen minutes later than Betelgeuse it sets nearly two hours earlier. This is because it is farther south than Betelgeuse and has a shorter path over which to travel from rising to setting. Betelgeuse is above the horizon for something more than thirteen hours, while Rigel is in view only about eleven.



When on the meridian, Rigel lacks four degrees of being half-way up to the zenith, and Betelgeuse lacks only four degrees of being two-thirds of the way up. The course across the heavens of Betelgeuse corresponds very nearly with that of Altair. The course of Rigel is about the same as that of Spica.

The other bright stars of Orion are hardly less interesting than Betelgeuse and Rigel. Bellatrix, which marks the left shoulder of the giant, is of the second magnitude, but still a very brilliant star. Three dazzling second-magnitude stars form the belt, or girdle, of Orion and lie half-way between Betelgeuse and Rigel. These are the stars supposed to be referred to by Job as "the bands of Orion." They lie in a straight line exactly three degrees long, and are rendered so conspicuous by their position and their intrinsic beauty that they serve as an aid in finding the constellation. They also provide a measure for estimating distances in this section of the skies. The upper one of them lies exactly on the celestial equator, and is a fine double star discovered by means of the spectroscope.

These three stars are called in some places the "three kings," and the line formed by them is sometimes called the "ell and yard" on account of its being just three degrees long.

Hanging from the belt is a curved line of stars known as the sword of Orion. The central star of the sword is distinctly hazy in appearance, and is no other than the famous Orion Nebula. It is one of the finest nebulae in the heavens and one of the few that can be seen with the naked eye. The one star that we see in it resolves through the telescope into four beautiful stars which form a figure sometimes referred to as the "Orion trapezium."

The fourth star, Kappa, which with Betelgeuse, Rigel, and Bellatrix forms the large parallelogram that outlines Orion's body is called Saiph and is situated in the right knee of the mighty hunter. Bellatrix signifies "Female Warrior" or "Amazon Star." Saiph means "Sword," although this star lies at some distance from Orion's pendent dagger.

With all its wonders and its beauties, it is not strange that Orion should be one of the most familiar and most admired of all the constellations. It is in the center of the galaxy that marches in brilliant procession across the winter skies. We watch for it between nine and ten o'clock in the evening late

in October, and our first view is of the curved line of faint twinkling stars that outline the left arm and the shield of lion's skin. Then one jewel after another emerges from the storehouse below the horizon until the whole splendid figure is before us. Its arrival is an announcement that the out-door season is past, that the nights are becoming more and more frosty, and that the gorgeous tapestry with which the autumn hills seem covered will soon fall away and give place to the lovely low tones of winter. We remember then that the juncos came that day and that as we crossed the coppice about dusk we heard them twittering cheerfully to each other as they settled for the night among the low bushes. And we make a mental note to see to-morrow if the little brown screech owl has moved into his winter quarters in the old catalpa-tree across the road.

During November and December, Orion becomes a familiar figure in the evening skies, rising a little earlier each evening until during the week on each side of New Year's Day he appears in the east just as daylight fades. In February he stands erect in the southern sky at nine o'clock in the evening. In March and April he is rushing furiously down the western sky after Taurus and the Pleiades, and by the middle of May he has disappeared from the evening view. In the latter half of July the very early riser can catch a glimpse of him in the east just before daylight. It was, no doubt, in this month that the "rosy-fingered dawn" took him for her lover, as she is reputed to have done.

.

## XIII

### The Heavenly Twins

ON the old astrological maps the constellation Gemini, or The Twins, was represented by two lusty-looking youths standing in an affectionate attitude with their heads practically consolidated, and each wearing a star as a sort of monocle in the outer eye. These stars are called Castor and Pollux, after the brothers from whose eyes they are supposed to shine. They are the brightest stars in the constellation Gemini, which is outlined by two almost parallel rows of stars beginning at Orion, running northeast about twenty degrees, and ending with Castor and Pollux.

Pollux is the brighter of the two and is the only first-magnitude star in the constellation. But it is almost as difficult to consider the two heavenly bodies apart from each other as it was to separate the two fond brothers for whom the stars were named.

The stars lie only about four and a half degrees apart and are so conspicuously a pair that each serves as an identifying mark for the other. Although 300 years ago Castor was reputed to be brighter than Pollux, this is probably not so. Nevertheless, Castor has some points of superior interest, as will be seen later on, and is entitled to be considered with respect and admiration in the gay company of our friends among the stars.

Castor and Pollux enter the eastern gate of the skies at about the same time that Betelgeuse and Rigel appear, but the Twins are so much farther north that there is no danger of confusing them with the two brilliant stars of Orion.

Betelgeuse and Castor come into view at almost the same moment, the former only about seven and the latter a little more than thirty-two degrees north of the exact east. About ten or twelve minutes later Rigel appears at a point a little more than eight degrees south of east, and after an interval of from ten to

fifteen minutes, Pollux shines out above the horizon about thirty-seven degrees farther north.

In November, when the twin stars first become a prominent feature of the early evening sky, Castor is leading. When they come to the meridian, about eight hours later, Pollux, being farther south and thus having a shorter course to run, has so gained on his companion that he reaches the meridian only about ten minutes later. At this point, which is their highest, Castor is nine-tenths and Pollux is about seven-eighths of the way up from the horizon to the zenith.

As they descend the western sky Pollux makes a still further gain, and in the latter half of the course they are seen side by side. They are particularly attractive during all of May and the first half of June, when they hang during the evening in the northwestern heavens shining out gently and kindly from the glowing background that the sun has left behind it.

Castor and Pollux were supposed by the ancients to exert a favorable influence over navigation, and the figureheads of ships often represented them. The Apostle Paul, when shipwrecked on the island of Melita, sailed from it "in a ship of Alexandria . . . whose sign was Castor and Pollux," and, of course, had a prosperous voyage to his friends in Rome.

The constellation Gemini is the third spring sign of the zodiac, and it is easy to see how the mere beauty of its chief stars, Castor and Pollux, may have fastened upon it the reputation of responsibility for the beautiful weather that comes early in June. At this season of the year position, atmosphere, and surroundings all combine to enhance the beauty and accentuate the individuality of these two beautiful stars. All, save one, of the radiant stars that have absorbed attention in this region of the sky during the winter and early spring have disappeared. The rainy Hyades, including Aldebaran, were the first to go. Stormy Orion followed soon after, and the ill-omened Dog Star went with him. And now in a comparatively starless environment the twin stars, beloved of sailors, dominate the western sky and shine side by side like two eyes benignly set to keep a protecting watch upon the world. It is not the sailor alone whose fancy is pleased with the kindly vigil they seem to keep. A landsman, too, may have pleasanter dreams if he will but take a peep through the west-

ern window and exchange friendly glances with them before settling down for the night.

Inasmuch as Castor and Pollux are so closely associated in our minds, in their times and seasons and their apparent position in the sky, it may be something of a shock to learn that they are in reality quite independent of each other and that they are going each his own way, in entirely different directions, towards an entirely different goal.

Castor is the farther of the two and is not very distant as stellar distances go. Its light comes to us in around 47 years and is about three million times farther from us than the sun. It requires about thirty-three years for the light of Pollux to reach us and the star is more than two million times as far from us as the sun.

Castor and Pollux are not only widely separated, but they are stars of very different spectral types. Pollux is yellow in color and similar to the sun and Capella, whereas Castor belongs to the large class of stars that includes Vega, Deneb, and about half of all of the brighter stars in the sky. They are usually white, though sometimes tinged with blue, as in the case of Vega. Some observers say that Castor is tinged with green, although stellar colors are probably not quite so finely differentiated.

Though the bond between Castor and Pollux is more apparent than real, Castor, at least, has a true companion. The companion is but a fraction more than one magnitude less bright than Castor itself and is greenish-yellow in color. It is bright enough to be seen with the naked eye, but we cannot separate the two stars without the aid of a small telescope because they lie so close together. The distance between them is only about four seconds, or one nine-hundredth of a degree.

The two stars revolve about each other very slowly, and though they were found nearly two hundred years ago to compose a double system of stars, they have not yet completed one full revolution, and their period is not exactly known.

Both of the principal stars composing Castor have been discovered to be spectroscopic double stars. In addition to these four stars, there is another small star near them that forms part of the system; this faint member is again a double; and so Castor is composed of six stars.

To swear by Castor and Pollux was common practice in an-

cient times, and later the name of the constellation itself came to be used in the same way. In Elizabethan literature we not infrequently find the expression "Oh, Gemini!" which by the modern rustic is corrupted into a slightly different word and gives him the expletive "By Jiminy!"

## XIV

### The Two Dog Stars

A STRAIGHT line from Aldebaran running through the belt of Orion and extending beyond it towards the south-east about twenty degrees will point out the Great Dog Star, Sirius, by far the brightest star in the heavens, and, happily, so situated that it can be seen in all southern latitudes and in all northern latitudes south of Greenland. Next to Polaris it is, perhaps, the best known of all individual stars, and this in spite of its not being visible in the latitude of the United States before midnight during the months when people are most out-of-doors and most likely to observe it—from May to November.

Sirius is called the Dog Star because it is the chief star in the constellation of the Greater Dog, which follows at the heels of Orion in his journey across the skies. The constellation gives the name of "dog-days" to the hot and trying period of the summer, including most of July and the first half of August. During this time Sirius rises just after the sun, and following it over the heavens each day, the star was supposed by the ancients to be in some way responsible for the malign heat of the midsummer sun and to exert an evil influence on the health of those exposed to its burning rays. Even now we frequently hear that dogs are likely to "go mad" at this season because "the Dog Star is in the ascendant." Indeed, the name Sirius comes directly from a Greek word signifying sparkling or scorching.

From Homer down to the present day few great poets whose themes were of out-door things have failed to refer to Sirius. A beautifully compact and comprehensive description of the star and its characteristics is given incidentally in the *Iliad*, where Achilles, mad with rage, was first seen by Priam "as he sped across the plain, blazing as the star that cometh forth at harvest-time, and plain seen his ways shine forth amid the host of stars

in the darkness of the night, the star whose name men call Orion's dog. Brightest of all is he, yet for an evil sign is he set, and bringeth much fever upon hapless men."

Sirius is the farthest south of any of the bright stars of winter that can be seen in this latitude, and in the sweep over its comparatively small arc in our skies it rises only about fourteen degrees higher than Fomalhaut, the most southern of the important autumn stars, and ten degrees higher than Antares, the most southerly first-magnitude star of summer. When Sirius is on the meridian, or at the highest point in its diurnal course from rising to setting, it is only about one-third of the way up to the zenith. It reaches this point about five hours after rising, and, hence, is above the horizon only ten hours out of each twenty-four.

We begin to look for Sirius in the evening sky about the middle of November, at which time it rises about ten o'clock. The real activity of the year is over. No flowers are in bloom save that here and there in some sheltered, sunny spot in the coppice a belated aster is cheerfully swinging its bright blue-starred branches, or some other flower whose season is long past has been enticed by the warm sunshine into showing yet one more blossom. To be sure, the witch-hazel, the very last flower of the year, which thriftily puts forth in the late autumn the bloom for next year's fruit, hangs full of the curious, straggling golden threads that constitute its flowers; but it stands bare of leaves and its empty seed-cups are open-mouthed as if with wonder and admiration at the success of the explosions by means of which it has lately peppered the neighborhood with its fruit. The heavy frosts have finished the work of painting the hills; the rich brown leaves are fast piling their soft winter covering over the yet warm earth; the smooth clean juncos have been flirting around in merry groups for three or four weeks, and the lovely Indian summer haze hangs like a soft mantle of gauze between us and the half-denuded trees.

In the skies all of the other bright winter stars have arrived, and some of the brightest of the spring and summer stars still linger to grace the company, when the king of stars arrives. No less than ten of our sixteen brightest stars are in view when the dazzling face of Sirius appears. On the eastern side of the heavens are golden Capella, rose-tinted Aldebaran, red-faced Betel-



geuse, fair Rigel, the gentle Pollux, and Procyon, the smaller Dog Star, all fresh and joyous and eager. In the southwest is Fomalhaut, serene and indifferent to what is about to transpire. In the west is Altair, just waiting above the horizon to witness the event and then drop out of sight. In the northeast are Vega and Deneb, bright and gay and twinkling with almost as many hues as Sirius himself.

He comes rich in many colors, twinkling fast and changing with each motion from tints of ruby to sapphire and emerald and amethyst. As he rises higher and higher in the sky he gains composure and his beams now sparkle like the most brilliant diamond—not a pure white, but slightly tinged with iridescence. It is worth enduring the chilliness of a clear, mid-November night at ten o'clock to see the assemblage of these eleven noble stars with their attendants. For mere brilliancy the celestial view is surpassed at no other season of the year.

Whereas Sirius is now white in color, some early references have indicated that a few thousand years ago it might have been red. Study of these passages, however, indicates that the proper translation is "fiery" rather than "red." It could scarcely be possible that a few thousand years, though a long time in the history of mankind, could possibly have led to major changes in the physical characteristics of a star like Sirius.

On Thanksgiving evening, which so often marks the first cold "snap" of the season, Sirius rises about nine o'clock. On Christmas evening it becomes visible a little after seven, and on New Year's about half an hour earlier. About the twenty-eighth of January it rises just as the sun sets, not far from five o'clock.

Being so low in the sky, it is easily observed at any point in its course. Except for the first thrill of pleasure that it gives when it appears in the late autumn, Sirius is never more attractive than during March and April. It is then at its highest during the early evening and shines brilliantly in the southwest. It sets with the sun about the twentieth of May, and for some days before that it has been invisible, its light being lost in the rays of our own great orb.

Sirius is one of the stars that we know to be comparatively near to us. Light travels from it to us in about eight years and a half, which means that it is about fifty-one trillions of miles distant, or more than five hundred thousand times farther away

than the sun. We get from the star only one seven-billionth of the light that the sun gives us; but Sirius in reality gives out about forty times as much light as the sun. It is a very gaseous and expanded body, however, and though very much larger and more brilliant than the sun, is comparatively light in weight, its mass exceeding that of the sun only about three times.

Sirius has a very interesting companion, which revolves around it in a period of fifty years. Its orbit is such that the two stars are nearer together at one time than at another, and it is only when they are farthest apart that the companion can be seen, and even then a strong telescope is necessary. When at their nearest, the companion is so completely lost in the brilliant light of Sirius that it cannot be found at all. The existence of such a companion had been long suspected, but it was not actually discovered until 1862, a year when the distance between the stars was increasing.

The faint companion is white in color, a fair match to Sirius. From this fact one deduces that their surface temperatures are about the same and that the stars have identical surface brightnesses. Since Sirius gives about 3,000 times as much light as its companion, we must further conclude that the companion is very tiny, scarcely larger than our earth. The mass of the fainter star, which is about half that of Sirius, is thus compressed into a very small volume. A teaspoonful of the highly condensed matter of such a white dwarf would weigh a ton. Such compression is characteristic of white dwarfs, which are believed to represent stars in the final stage of their history.

The other Dog Star, Procyon, lies in the constellation of the Little Dog (*Canis Minor*). To locate Procyon, run a line a little more than twenty degrees northeast from Sirius, or about the same number of degrees east from Betelgeuse. Procyon, Sirius, and Betelgeuse are at the angles of an almost equal-sided triangle, and the same three stars form with Rigel a figure that is almost a square. Procyon is twenty-three degrees directly south of Pollux and forms a triangle on the north with Pollux and Betelgeuse of just about the same size as the triangle it forms with Betelgeuse and Sirius on the south. It rises an hour and a half later, and sets an hour and a half earlier, than Pollux.

The name Procyon means "before the dog" or, more literally, "precursor of the dog." The star received this name because it

appears in the east from fifteen to twenty minutes before the Great Dog Star. It rises very little north of the exact east, reaches the meridian about six hours and twenty minutes later, and sets between two and three hours after Sirius. At its highest point it reaches about three-fifths of the way up to the zenith.

Procyon suffers somewhat in reputation by reason of its connection with Sirius. The fact that it is known as "the other Dog Star" and "the Little Dog Star" indicates inferiority of some sort. It is, in fact, a most beautiful star, and is only the sixth in order of brightness among the stars seen in our latitude. It is very distinctly individual, being the only one among the beautiful group of winter stars that is lightly tinged with yellow. Procyon is definitely cooler than Sirius and in some ways resembles the sun, though it is considerably brighter.

Procyon is also one of the near-by stars, but it is not so near as Sirius. Light travels from it to us in a little more than eleven years; so that the star is about seventy trillions of miles away from us. It gives out six times as much light as the sun, and is in no sense an insignificant body or one to be known only as announcing the imminent arrival of Sirius.

Procyon also has a companion of its own, a yellow-white star of about the eleventh magnitude, which revolves around it in about 40 years. It is a matter of coincidence that, like the companion to Sirius, the body accompanying Procyon is also a white dwarf. In other words, the faint companions to both of the dog stars are the enormously dense white dwarfs. The companion of Procyon was not discovered until 1896.


## XV

### Regulus

THERE is no single line by which Regulus can be exactly pointed out to one who does not know the constellations. Still it can be easily found. A line forty degrees long drawn through the pointers backward—that is, away from the North Star—will end just a little to the west of a conspicuous group of stars known as the Sickle (and really resembling that instrument), and Regulus is the brilliant jewel that sparkles at the end of the handle.

In June and July, when Antares and Spica are both shining, Regulus may be found by a line drawn from Antares through Spica and extended about fifty degrees. The line will unmistakably indicate Regulus, but the star will be well down in the west and not far from setting. When Arcturus is visible and not too far down in the west, Regulus may be found at a point about sixty degrees west (a very little north of west) of Arcturus. A triangle is formed by Spica, Arcturus, and Regulus, with Regulus at the acute angle pointing towards the west.

Among the brightest stars, Regulus comes and goes rather between seasons. It is not strictly a summer star or a winter star, though it may be seen in both seasons. It is, in fact, visible at some time between sunset and midnight for eight months in the year. It arrives after the galaxy of brilliant winter stars are in full swing across the heavens and before the gay company that gladdens our view in the spring has begun to assemble. It appears above the horizon a very little north of east about nine o'clock in the evening of New Year's Day. One may even have seen it a little later in the evening than this some days before Christmas; but Regulus does not have the insistent quality of brightness that commands the regard of the ordinary observer



THE SICKLE

until it has reached a position and an hour when one can observe it conveniently.

From the time it first attracts our attention until Arcturus and Vega come, Regulus is the chief ornament of the eastern skies and shines like a veritable diamond of the highest quality, fit to be set in the jewelled hilt of sword or scimitar of fine Damascus steel rather than the handle of a homely garden implement. It is a singularly clear, white, sparkling star, with just a touch of blue. But in spite of its brilliancy it seems a much smaller point of light than many of the bright stars. It is next below Deneb and Beta Crucis in its rank among the stars as to brightness, and the stars are so nearly equal that some astronomers have given Regulus the highest place. Modern photometry, however, ranks it sixteenth in order of brightness among the stars visible in the northern hemisphere. It ranks twenty-first among all the first-magnitude stars.

It requires about six hours and three-quarters for Regulus to reach the meridian, and it is then about one degree more than

two-thirds of the way up to the zenith, or the point exactly over our heads. It shines all night in January, February, March, and April, rising in February just as the sun sets. In May, June, and July it may be seen travelling towards the western horizon during the first half of the night. By August it is so near the sun that even with a favoring atmosphere we can barely get a glimpse of the star during the first few days of the month, as it falls below the horizon close after the sun; and it remains in this region most of September. In the early morning hours of October and November it may be seen gradually gaining on the hours until it comes back to its familiar place in the east.

Blue-white Regulus is a trifle hotter than Vega, Deneb, or Sirius, but not quite as hot as Spica. It certainly ranks among the stars having high surface brightness. Regulus is about 140 times brighter than our sun, and its distance from the solar system is so great that we do not receive the light from the star for some 77 years after it has started toward us, moving though it does at the enormous rate of 6 trillions of miles a year.

Regulus is in the constellation called Leo, or the Lion. The blade of the Sickle marks the shoulder of the Lion, and Regulus lies at its heart. The star is sometimes called "Cor Leonis," meaning "the Heart of the Lion." The name Regulus means "chief."

The constellation Leo is one of those that the sun passes through in its apparent annual path across the heavens. The sun stays in this constellation a little more than five weeks, entering it at the beginning of the second week in August. Two weeks later the sun passes Regulus, and as the star lies almost exactly in the path of the sun, they would seem to an observer on the earth to almost touch each other. But at this time the light of Regulus is lost in the sun's rays, and we do not see this interesting meeting.

Another interesting star in Leo, though not of the first magnitude, is Denebola. It marks the tuft at the end of the Lion's tail, and its name means "the Tail of the Lion." It lies about twenty-five degrees east of Regulus and a very little to the north of it. It is the brightest star in a small triangle which a glance towards the region indicated will reveal. A small telescope shows Denebola as a beautiful double star of contrasting colors. The pair does not form a physical system, however. The stars do not

revolve around one another but lie merely in the same direction.

Within the curve of the blade of the Sickle is the point from which the meteors that come along in the middle of November seem to radiate. They are on this account called "Leonids."

## XVI

### The Number of the Stars

ON a clear starlit night the sky seems so thickly set with stars that, on first view, it is difficult to believe that within two hours one could count every one of them. Yet this is true, for to the ordinary eye there are not more than a thousand stars visible at one time. The very keenest eye without mechanical assistance cannot discern in the heavens, north and south, more than seven thousand, and barely five thousand are visible to the ordinary sight. Of this entire number nearly one-fourth are too far south to be seen at all in our latitude; and of those that can be seen here only one-half are visible at once, the others being below the horizon and visible only in the other hemisphere. And, finally, of those that are really above the horizon and could otherwise be seen, the bright stars are somewhat obscured by the dense atmosphere while near the horizon, and the faint ones are completely blotted out for a considerable distance farther up. This is why there seem to be no faint stars low down in the sky. In reality they very largely outnumber the bright ones in every part of the heavens, but they can be seen only when they rise high enough to escape the heavy lower atmosphere.

The stars that we can see with the naked eye (astronomers call them *lucid stars*) are, of course, a very small proportion of the total number of stars in the universe. As to the total number of stars within our galaxy, astronomers generally believe that there must be approximately 100 billion. It is estimated that the largest telescope will reveal approximately 2 billion. Remember, these are just the stars of our own galaxy. When we realize that there may be as many as a trillion galaxies accessible with large telescopes, each containing perhaps as many stars as our own galaxy, the total number of stars proves to be tremendous.

The difference of opinion among astronomers as to the limi-



tation of the stars is based in part upon their ideas of the destructibility of light. It is certain that we do not get an infinite amount of starlight. If we did, all space, they say, would be filled with a diffused glow and there would be no such thing as darkness. For it is not necessary that we should actually see the stars individually in order to have light from them. Those that we can see with the naked eye give only about ninety per cent of the total starlight received by us. The naked-eye and the telescopic stars combined give only about one one-hundred-and-eightieth as much light as the full moon. But on a bright starlight night the light we receive is equal to about one-sixtieth of full moonlight, which shows that we get three times as much light from the stars whose rays are too faint to be seen as from all the other stars together.

It has been shown by photography that we get light from stars that are invisible to the eye in even the largest telescope. The photographic plate has an advantage over the eye in that the plate can store light as the result of a long exposure. The eye, on the other hand, has no storage capacity. In fact, because of fatigue, the eye, used continuously over a period of time, usually sees less at the end of that interval. The faint stars, so recorded, which will ever be invisible to the less sensitive human eye, are nevertheless contributing light to the sky, for they would make no impression on the photographic plate if their light did not reach it.

Observations have shown that interstellar space contains a certain amount of fine dust particles. The amount of dust in any given volume of space is relatively small, however. If we used a vacuum cleaner to sweep out a volume as large as our earth, for example, of even the dustiest regions, we should collect at most about a cupful of dust. Even so, distances in interstellar space are so vast that, in certain regions, the dust appreciably dims the stars. Our Milky Way would be much brighter if dust clouds did not obscure the stars behind them.

Einstein's theory of relativity has indicated that space is not infinite. At great distances, a natural curvature of space tends to limit the volume. Hence, the number of stars, either in our own galaxy or in other galaxies, cannot be infinite.

There are certain bodies in space which, for want of a better name, we speak of as dark stars: that is, stars that give no appre-

cial light. They make themselves known in some instances by passing at regular intervals between us and a bright star, thus showing that they are revolving around it. In other cases they make the star swerve from what its course would naturally be, thus showing the influence of some invisible body. Most of these so-called dark stars, however, are not completely dark. But there well may be many dark bodies in space, either dead stars or cold matter like our planets or meteors. But at present they need not be taken into account.

The telescope and camera both give evidence of a limit at least to the number of the stars that we can really know. As telescopes were made larger and larger a greater number of stars became visible; and it seemed as if the number we could see depended solely upon the power of the glass. It has been found, however, that beyond a certain point the number we can see does not increase in proportion to the increased size of the telescope. This indicates that we must be approaching the limit of the number of stars that send their light to us. The same thing is shown by the camera. After a certain length of exposure, additional time will add no more than a few stars to the number already on the plate.

There are about twenty well marked degrees of brightness in the stars seen through the largest telescopes. For the first ten or twelve of these the number of stars in each grade is about three times greater than in the grade preceding it in order of brightness. That is, we have of the first degree of brightness twenty-one stars; of the second degree, about sixty; of the third, one hundred and eighty, and so on. But after about the twelfth degree of brightness the number of stars does not increase in the same proportion; the increase as shown by both the telescope and the camera is comparatively small.

It is not to be supposed that, because telescopes and the camera have not yet revealed them, astronomers think there are no more stars to be discovered. But the evidence of the telescope and photography shows we are approaching that limit. The exact number will probably never be known. Certainly it will never be known by means of the telescope; for we seem to be reaching a point where it will be physically impossible to make larger telescopes that can be effectively used. The last extension of our knowledge of the number of the stars will probably be

reached only indirectly. For, even if we cannot see the stars individually, their collective gravitational effects may be sufficient to influence the motion of visible stars in that region of the sky. The more rapidly the stars move the greater the mass of these invisible stars. But even by this method we cannot infer the existence of truly enormous quantities of invisible, undiscovered, dark matter.

What is not recorded on the photographic plates of the future will remain to us unknown. We can be reasonably sure of only one thing: that, in view of the evidence we have of the decrease in the number of the stars as they grow fainter to us, the ones remaining undiscovered are practically but a few stragglers along the "fringes of the visible universe"—few, even though there may be really many millions of them.

While the number of stars that are known is increasing somewhat all the time, astronomers are not so much trying to make discoveries of larger numbers as to learn something more about the stars that are already known. The number now classified and catalogued is something near eight hundred thousand.

A great international photographic chart of the heavens, completed with international cooperation, showed more than 40,000,000 stars. And the greatest survey of all, a series of photographs taken with a powerful Schmidt camera at Palomar Observatory, has revealed so many stars that no accurate census of their number is available. Unfortunately, this Palomar survey covers only the stars visible from the northern hemisphere. A similar survey for stars in the vicinity of the south celestial pole will probably be completed in the next few years.

.

## XVII

### The Names of the Stars

**H**APPILY for the interest of the ordinary observer, nearly all the brighter stars, and a few of the fainter ones, have individual names. Sixty or more of these names are in common use even among professional astronomers. Most of these belong to the so-called navigational stars, the stars that navigators observe to find their positions on the sea or in the air. A few more names are well known and are used occasionally, and a number which are now practically obsolete appear in old catalogues, or historical lists, or in some piece of ancient literature. Less than a hundred names will cover all the stars of the first and second magnitudes; and besides these there are not many that have an individual interest for an observer without a telescope. When one learns to know fifty or sixty stars so that he can always recognize them and call them by name, he begins to feel that he has a pretty large acquaintance in the heavens; and the knowing of other individuals that may chance to interest him is then the mere matter of an introduction.

Most of the individual stellar names originated with the ancient astronomers and have a meaning that describes some characteristic of the star, some trait, or color, or position in the constellation. All of them were probably descriptive at first, but the original meaning is not always clear. Sometimes designation was made by a circumlocution such as "the Heart of the Lion," "the Eye of the Bull," "the Foot of the Giant." This was, of course, too cumbersome and limited a method to answer the needs of modern astronomy. But many Latin and Arabic names are still used which in one word describe the place the star holds in its constellation; as, for instance, Achernar, meaning "the End of the River," and Denebola, "the Tail of the Lion."

Very early in the seventeenth century, astronomers adopted

the plan of designating the stars in each constellation by the letters of the Greek alphabet and adding the name of the constellation in its Latin form in the genitive (or possessive) case, thus making the name of the constellation a sort of family name for all the stars composing it, and the Greek letters the given names. Thus in the constellation Taurus we have Alpha Tauri, meaning Alpha of Taurus; Beta Tauri, Gamma Tauri, and so on. When the Greek alphabet was exhausted, the Roman letters were used. Beginning with alpha, the stars of each constellation were named in the order of their brightness. In the main, though, when several stars in the constellation chanced to be of nearly the same degree of brightness, there seems to have been no nice distinction made between them. A major exception is the Big Dipper, whose seven bright stars carry the Greek letters from Alpha to Eta in order from bowl to handle.

A century later another catalogue was made, and in this numbers were used instead of letters for the given names, the genitive case of the constellation still being used as a family name. Several special catalogues have been made in which also the stars are indicated by numbers, such as Lalande's, the British Association Catalog, the Harvard Photometry, and the Henry Draper Catalog. In such cases the reference is simply by number and name of catalogue. Some stars that have in a way become famous by reason of some individual peculiarity or characteristic still have no designation except by number. Among them, and spoken of more particularly elsewhere in this book, are 61 Cygni, 1830 Groombridge, and some others. Stars are also designated by their position in the heavens, which is expressed in the terms of astronomical measurements, right ascension and declination, the celestial analogs of terrestrial longitude and latitude.

A result of these various methods of naming the stars is that nearly every bright star is designated in several different ways: by its individual name, its Greek letter name, its number in the various catalogues, and its position in the heavens. But the ordinary observer need not much trouble himself with learning them all. One can have an intimate knowledge of nearly all the naked-eye stars without once resorting to figures for names, or without even knowing what right ascension and declination are. Astronomers have evolved these several schemes of designation only for their own special needs. Rarely has a star known only by its num-

ber become, for any reason, of general interest. There are not now more than half a dozen such stars. Probably 61 Cygni (that is, the star numbered 61 in the constellation of the Swan, or Cygnus) is the best known of these, because it is the first star whose distance from us was measured. Despite its nearness, however, it is of magnitude 5.5, one of the fainter naked-eye stars.

The method of designation most common among writers on astronomy for the brighter stars (when the individual name is not given) is that of the Greek letter joined to the name of the constellation. After the constellations are once known, this is an easy method, and, even for the non-professional observer, has several advantages. It not only names the star, but assigns it to a constellation, and gives an approximate idea of its degree of brightness as compared with other stars, and particularly as compared with other stars in the same constellation. Alpha of a constellation is almost always the brightest star in the group. The exceptions to this are few and mainly accidental. One instance is that of Castor and Pollux, the two bright stars of the constellation Gemini, or the Twins. Pollux is the brighter of the two, yet Castor is Alpha Geminorum and Pollux is Beta Geminorum.

It is not often that the average reader will care, or need, to know the names of more than the seven or eight brightest stars of a constellation, the ones designated by the first seven or eight letters of the Greek alphabet: alpha ( $\alpha$ ), beta ( $\beta$ ), gamma ( $\gamma$ ), delta ( $\delta$ ), epsilon ( $\epsilon$ ), zeta ( $\zeta$ ), eta ( $\eta$ ), and theta ( $\theta$ ). Rarely, indeed, will he care to know even that many. In the Great Bear (*Ursa Major*) the seven brightest stars, the ones that form the well-known Dipper, are of almost equal importance and brilliancy. But this is an exceptional case, and more often only three or four stars in a constellation, the alpha, beta, gamma, and delta of the group, have any special interest. Occasionally one of the fainter stars calls for particular attention, as in the case of Mira, the Wonderful, in the constellation of the Whale, or Cetus, which is designated by the fifteenth letter of the Greek alphabet, omicron, thus:  $\omicron$  Ceti.

The Greek letter method of naming the stars has, then, its conveniences. But, after all, when a star, like anything else, comes to have any special individuality for us we desire a name for it that is all its own. When we come really to know splendid Arcturus, we do not care to call him Alpha Boötis any more than

we want to call a dainty spring beauty *Claytonia virginica*, or the golden-eyed white daisy *Chrysanthemum Leucanthemum*. And, happily, we are not often put to this discomfort, for there is no star of the first magnitude in the northern hemisphere that has not an individual name. In the southern hemisphere, however, Alpha Centauri, Beta Centauri, Alpha Crucis, and Beta Crucis are more commonly referred to by their Greek-letter designations than by their proper names, which are Rigil Kentaurus (or simply Rigil Kent), Hadar, Acrux, and Becrux, respectively. The last two names are specially coined to designate the stars for navigational purposes. Alpha Centauri, we have seen, is the star that is nearest to us.

I remember feeling distinctly grateful to the old astronomers for having given individual names to much less conspicuous or important stars. A window in my bedroom opens out towards the northeast, and during the autumn and early winter the Dipper is visible there a good part of the night. Many times it has happened that I have suddenly opened my eyes in the night and found there beaming down upon me the soft rays from the star Mizar, which lies just in the bend of the handle of the Dipper and next to the end star, and its tiny companion Alcor, that seems to send its own little beams almost along the same path. The simple friendliness that I felt in encountering them in this way came as much, I think, from my being able to greet them by their own names as from their intrinsic beauty and steadfast cheerfulness.

## XVIII

### The Light of the Stars

STARS are classified according to their brilliancy into magnitudes; and as applied to them, the word magnitude has a specific meaning. It has reference solely to their brightness and has nothing to do with their proportions. Many stars that appear to us very small and faint we know to be of immense size, far exceeding some of the most resplendent stars of the heavens; so there is no simple relation between the real and apparent magnitude of a star—that is, between its size and its brilliancy.

This use of the word magnitude was fixed in astronomy in the early days of the science, before the telescope had been invented and before there was any way of judging of the dimensions of a star other than by its appearance to the naked eye. The stars that seemed the largest and brightest were supposed really to be so, and were said to be of the first magnitude; the next brightest were called second magnitude stars, the next third magnitude, and so on. All the stars visible to the naked eye were divided according to their lucidity into six magnitudes, and modern astronomers have found it convenient to continue this way of describing their comparative luminous power, and have applied the same principle in describing telescopic stars even below the twentieth magnitude.

It is, of course, an arbitrary classification, for there are stars of every degree of brightness between Sirius, the brightest star in the heavens, and the faintest star we can see; and, so far as accuracy is concerned, we might as well have had sixty magnitudes among the visible stars as six. Moreover, with no more accurate measurement than one can make with his eye, it is not easy for the ordinary observer to tell the difference between the amount of light we receive from a faint star of one magnitude and a bright one of the next lower magnitude. Since naked-eye observation was the only way of determining stellar magnitudes



among the ancient astronomers, there was considerable difference of opinion concerning the brightness of such stars as were on the border lines of each magnitude. To the first magnitude there were assigned anywhere from fourteen to twenty stars, and in the lower magnitudes the difficulty of agreement was even greater.

Modern astronomers, while retaining the old method of classification, have endeavored to systematize it by the accurate measurement of the amount of light we receive from each star. This is done with an instrument known as a photometer, by means of which very slight gradations in the light of the stars can be determined. With this instrument practically all the naked-eye stars have been measured, and they have been catalogued accordingly. Modern photoelectric photometers are extremely accurate and sensitive. They are capable of recording gradations of brightness as small as one one-thousandth of a magnitude.

In making the measurements, the astronomer adopts some standard of light, compares the light of each star with that standard and then determines the stellar brightness in terms of the basic scale of magnitudes. In the Harvard Photometry, the Pole Star (Polaris) was originally the adopted standard star taken for comparison. It is a conspicuous and important star that everyone knows, and has long been rated as of the second magnitude—even in the days of rough judging of the brightness of stars. Unfortunately, however, Polaris is slightly variable. To extend the photometric scale to fainter magnitudes and to provide standards of comparison that would always be visible from the northern hemisphere, astronomers have measured very carefully the brightness of a number of stars in the vicinity of Polaris. These form the North Polar Sequence. Now, when an astronomer wants to determine the brightness of some particular star in the sky, he measures it very carefully with a photometer and then turns to the North Polar Sequence for checking and standardizing his measures.

One would think that the determination of stellar magnitudes would have been based on either the brightest star or one of the brightest as a standard and that all other stars would have been graded from that. This is probably what would have been done if the stars had not been classified as to brightness until the

delicate instruments for measuring their light were invented. As it is, the classification was made many hundred years ago, and though modern astronomers are endeavoring to express the comparative gradations of light as accurately as possible, they still do not much disturb the distinctions that have long been familiar.

By photometric measurements the gradations of light between the different magnitudes are expressed in tenths by decimal fractions. Thus the magnitude of one star is 2, that of another a little less bright is 2.1, and so on to 2.9, which is almost as faint as a star whose magnitude is 3. The larger the fraction the nearer the magnitude of the star is to the next magnitude below the one in which it is classed, and the higher the number of the magnitude the fainter the light of the star.

For the ordinary observer it is sufficient to know that a star is of the first, second, third, or whatever magnitude, as the case may be. These nice distinctions in tenths between degrees of brightness are useful more to the serious student than to the general reader. The usual way of classifying is to designate a star as of the magnitude to which it is nearest. Consequently each magnitude reaches out on each side and takes in all stars that are not more than five-tenths of a magnitude from it. For example, all are called first-magnitude stars whose magnitude is brighter than 1.5. If a star's brilliancy is found to be 1.6, it is classed as a second magnitude star, and all are of the second magnitude from this down to 2.5; and similarly the other magnitudes are reckoned.

This is true of all the magnitudes except the first. This has its limits only on one side. It includes all stars that are brighter than 1.5 magnitude. There are but twenty-one stars of the first magnitude, but the individual lustre of them varies so as to cover nearly three magnitudes. To overcome this difficulty astronomers express the magnitude brighter than 1 as 0, the magnitude next brighter as  $-1$ . The magnitude of Sirius, the most brilliant of all the stars, is  $-1.42$ . Alpha Crucis, Spica, and Antares, all of the southern hemisphere, are almost exactly first-magnitude stars. Five others are less bright than Antares, and twelve are between Alpha Crucis and Sirius.

According to the most recent photoelectric measurements, the visual magnitudes of the twenty-one brightest stars are as follows:

Sirius (Alpha Canis Majoris) .....	—1.42
Canopus (Alpha Carinæ) .....	—0.72
Alpha Centauri .....	—0.27
Arcturus (Alpha Boötis) .....	—0.06
Vega (Alpha Lyrae) .....	+0.04
Capella (Alpha Aurigæ) .....	0.05
Rigel (Beta Orionis) .....	0.14
Procyon (Alpha Canis Minoris) .....	0.38
Achernar (Alpha Eridani) .....	0.51
Beta Centauri .....	0.63
Altair (Alpha Aquilæ) .....	0.77
Betelgeuse (Alpha Orionis) .....	{ 0.1 max. 1.2 min.
Aldebaran (Alpha Tauri) .....	0.86
Alpha Crucis .....	0.9
Spica (Alpha Virginis) .....	0.91
Antares (Alpha Scorpis) .....	0.92
Pollux (Beta Geminorum) .....	1.16
Fomalhaut (Alpha Piscis Austrini) .....	1.19
Deneb (Alpha Cygni) .....	1.26
Beta Crucis .....	1.28
Regulus (Alpha Leonis) .....	1.36

The light ratio agreed upon between the different magnitudes is 2.5. That is, a standard star of each magnitude is two and a half times brighter than a standard star of the next lower magnitude. Or, to put it the other way, each magnitude is four-tenths as bright as the next higher magnitude. This makes a first-magnitude star one hundred times brighter than a sixth-magnitude star, and a sixth-magnitude star ten thousand times brighter than one of the sixteenth magnitude. In the same ratio, a first-magnitude star would be nearly four hundred thousand times brighter than a fifteenth-magnitude, and thirty-seven million times brighter than a twentieth-magnitude. This would make Sirius something over two hundred million times brighter than the faintest star of the twentieth magnitude.

But notwithstanding this vast superiority in brilliancy of the brightest stars, we get more starlight from the faint stars as a whole than from the bright ones as a whole. This is on account of the large number of the faint stars; for it has been found that the number of stars of a given magnitude is about three times greater than the number of stars of the next brighter magnitude. Hence, though a first-magnitude star is two and a half times brighter than a second-magnitude star, there are three times as many second-magnitude stars, and consequently we get more light from them as a whole. Though a first-magnitude star is

three thousand times brighter than one of the tenth magnitude, yet there are seventeen thousand times as many of the tenth magnitude as of the first. This progressive increase in number is an approximate estimate, but it is deduced from actual catalogues of the stars and is fairly accurate. As previously noted, however, the increase in number of three times between one magnitude and the next holds only for the brighter stars. The ratio decreases toward the fainter stars.

According to this ratio of progression, if one makes due allowance for the way in which one magnitude merges into another, the six or seven thousand stars visible to the naked eye are distributed roughly as follows:

21	stars	of	the	first	magnitude,
60	"	"	"	second	"
180	"	"	"	third	"
500	"	"	"	fourth	"
1500	"	"	"	fifth	"
4500	"	"	"	sixth	"

Thus the total number of stars visible to the naked eye is about 7000. How many stars one really sees of this seven thousand depends upon the state of the atmosphere and the keenness of one's sight. Smoke, dust, haze, or even the scattering of light by a transparent atmosphere greatly reduces the brightness of stars toward the horizon, for one is looking through increasing thicknesses of air. As a result, most persons under ordinary conditions fail to see nearly two thousand of them, and it takes an eye of remarkable power to see them all. However many one sees, there are always many more just on the verge of visibility that seem to come and go. This is one reason why there seem to be so many more stars in view than there really are.

It is to be remembered that all this applies only to the stars as we see them, and has no relation to their real size or their light-giving power. They may be brighter because they are nearer to us, or because they are really more brilliant and give out more light than others; but none of these qualities is determined by their apparent luster. As a matter of fact, some of the brightest stars are very well known to be not the largest.

On the average, of course, the brightest stars tend to be the nearest to us, though there are many exceptions. The converse

of this statement is far from true, however. The stars nearest to us, with a few exceptions, do not tend to be among the brightest. This paradox we can easily resolve. A given volume of space has in it only relatively few very luminous stars, but it has enormous numbers of stars whose luminosity is very low. For us to be able to see these intrinsically faint stars they must be extremely close to us. Thus, of the seventeen nearest stars, ten are too faint to be seen with the naked eye and one of them, Wolf 359, is of magnitude 13.5, a thousand times too faint to be seen without telescopic aid. The intrinsically bright stars, however, are intensely luminous and can be seen to enormous distances. Even though they are relatively rare in a given volume of space, the total volume through which they are visible is enormous. As a result, it is safe to say, that with a few exceptions, most of the stars visible to the naked eye are fairly distant suns. The best known exceptions to this rule include Alpha Centauri, the nearest star, Sirius, and Procyon.

## XIX

### The Distances of the Stars

THOUGH astronomers had been working for many years on the problem of determining the distances of the stars, it was not until 1838 that they succeeded in getting what seemed to be any fairly accurate measurements. In that year the distance of 61 Cygni was measured with reasonable correctness; and at about the same time it was found that Alpha Centauri was even nearer than 61 Cygni. Strangely enough, these two stars are still counted among our nearest neighbors. The faint companion of Alpha Centauri, of the eleventh magnitude, has proved to be a member of that system. It may be a trifle nearer to us than Alpha Centauri and hence has sometimes received the name Proxima. The assignment of such a name is tempting fate. But, curiously enough, the name may be justified. However, it is well within the realm of possibility that, among the millions of still fainter stars on our photographic plates, may be found one or more that lie still nearer. If such a star is discovered, of course, the name Proxima will then have to be completely discarded.

For many years 61 Cygni was supposed to be the next nearest star. However, in the meanwhile, at least seven stars have been found to lie closer than 61 Cygni, among them the brightest star in the sky, Sirius.

The distances of a number of stars have been determined since 1839; and through their knowledge of the distances of these, astronomers have been able to at least place limitations on the distances of others, and taking color, brightness, and other characteristics into consideration, have made inferences also as to size and constitution which it seems reasonable to believe are very near the truth. The ascertainment of distances has opened up a very large field of investigation; for, until the distance of a star is known, we can know but little of its size or

light-giving power, and hence little of its real nature, though its chemical constitution may be revealed by the spectroscope. Two stars may be apparently of equal brilliancy, but when we learn that one is many times farther from us than the other, we realize how much brighter the more distant one must be. Take, for instance, the two brightest stars, Sirius and Canopus. They are both so much brighter than a standard first-magnitude star that they are really in a class by themselves, though called first-magnitude stars. Sirius is the brighter of the two, and it is five hundred thousand times farther from us than the sun. Canopus is so far away that it successfully resisted, for a long time, all efforts of astronomers to measure its distance. Finally, special techniques solved the problem and it was then found that Canopus is indeed one of the most luminous of all of the first-magnitude stars.

Measuring the distance of a star is called by astronomers finding its parallax. Roughly speaking, parallax is the apparent change of position that an object seems to make when viewed from different points. If you hold a pencil out in front of you about one foot and look at it with first one eye and then the other, you will see that it seems to change its position relative to any object back of it. If you hold the pencil farther away, you will see that the change of position seems less, and if you place it still farther away, there seems to be no change of position at all. Then if you will move your head from side to side and thus lengthen the line between the two points of observation, which was at first only the distance between your eyes, the pencil will have to be placed still farther away before it seems to remain stationary when looked at from both points of view. The longer the distance between the two points of observation, the farther away the object observed will have to be to remain in apparently the same place when viewed from each point. When mathematicians know the length of the base line (that is, the line connecting the two points of observation), and can measure the angle made by the meeting of the other two lines (that is, the lines from the two points of observation to the object observed), they can tell how far away the object is.

When looked at in the ordinary way, a star never seems to change its position in the heavens. Even if viewed from points whose distance from each other is the whole diameter of the earth, eight thousand miles, it shows no change of position.

At first thought one would think that we could not find a line longer than that from one side of the earth to the other. But astronomers have found one in space. It is the diameter of the earth's orbit, or the path over which the earth travels in its yearly journey around the sun. This diameter is 186,000,000 miles long, or more than twenty-three thousand times as long as the diameter of the earth. Observations of the stars whose distance is to be measured are made when we are at one point in the earth's orbit, and then again six months later, when we are at the opposite point, or 186,000,000 miles removed from the first. A few stars change their apparent positions when thus observed from the two ends of this great line; but the changes are still very slight. By strict mathematical calculation we know, however, that any star showing a change of position of only one second, the smallest subdivision of a circle, when viewed from each end of a line 186,000,000 miles long, must be a little more than nineteen trillions of miles from us. Thus far no star in the heavens has been found to change its position this much; or, as astronomers say, no star has yielded a parallax as large as one second ( $1''$ ). Consequently, as far as we know, there is no star as near to us as nineteen trillions of miles. The largest parallax that has been found is .75 of a second. Actually, a minimum of three observations is needed to determine the parallax of a star. The third observation enables the astronomer to distinguish between the apparent drift of the star across the sky caused by the relative motion of star and sun and the shift caused by the parallax.

The sun is ninety-three millions of miles from us, yet if we were journeying in space, we might reach the sun and then go two hundred thousand times farther away and still not come near any fixed star. Indeed, we should have to go still eighty thousand times the sun's distance from the earth, for a parallax of .75 of a second means a distance of seven trillions of miles more than the nineteen trillions. In other words, the nearest star that has been discovered is about twenty-six trillions of miles away, which is equal to two hundred and eighty thousand times the distance of the earth from the sun.

Other stars are many times farther away than this. Their distances in miles expressed in figures are not much easier to comprehend than infinity. Most astronomical numbers are too large



to comprehend and too cumbersome for easy use. For this reason the distance of the earth from the sun (93,000,000 miles) has been adopted as the astronomical unit, and in expressing long distances we ordinarily say so many times the sun's distance. But some of the stars are millions of times the sun's distance, which still gives us an awkward lot of figures to deal with. Astronomers have, therefore, chosen another unit of length for expressing stellar distances. This unit is the distance over which light passes in a year. We know that light travels at the rate of 186,300 miles in a second. This equals about six trillions of miles in a year, or about sixty-three thousand times the sun's distance. Therefore, for every six trillions of miles that a star is distant we speak of it as one light year away. The nearest star, then, is a little more than four and one-third light years away, which means that its distance is more than four and one-third times six trillions of miles.

The light that we see from the nearest fixed star left there more than four years ago. From many stars we do not see it for a hundred years, or for many hundreds of years. From some of the faintest stars it may take light many thousands of years to reach us. Streams of light are travelling towards us now that set out on their journey long before Christ was born and have not yet reached us. The stars from which they come might no longer exist. The Pole Star, which we think is the most stable thing in the universe, might be extinguished for nearly half a century and we should still be ignorant of the fact; for the light that we now see probably started on its journey to us almost fifty years ago. And this though the velocity of light is so great as to send it around the entire circumference of the earth (25,000 miles) more than seven times in a second. From the nearest fixed star our sun, which gives us ninety billion times more light than a first-magnitude star, would shine only as a star of the second magnitude. Pluto, the farthestmost planet of the solar system, is about three billions of miles from the sun; yet between us and the nearest fixed star we could line up several thousand such solar systems just touching one another and still have some room left over.

We think sound travels fast when we say that it goes at the rate of one thousand and ninety feet per second. But if it were possible for sound to travel to us from the nearest fixed star, it

would, even at this rate of speed, spend three millions of years on the journey through space. Fortunately sound is dependent on our atmosphere for its motion and does not travel through space as light does. If it did, we would probably have such a mighty and continuous thunder that our ears would have to be constructed very differently from what they are now to stand the shock; for tremendous explosions and upheavals are going on all the time among the stars, of which our sun is one.

Even though the sound waves from such disturbances do not plague us, we have nonetheless become aware of the disturbances because of the radio noise such disturbances produce. During periods of intense sunspot activity, sensitive radio receivers pick up a violent hissing noise that swishes on different frequencies of radio waves. Many objects in the stellar system also send out a hissing noise indicating the presence of great disturbances. One of the chief sources of radio noise is the Crab Nebula, in Taurus. This object is a planetary nebula. It actually represents the remnants of a star, a supernova, that exploded in the year 1054. Ancient records show that a bright star appeared in this position in that year.

Measurement of stellar parallaxes is one of the most delicate of astronomical operations. In such observations allowance must be made for errors of various kinds. Even personal peculiarities must be taken into account; for, with the same honesty of purpose and working with the same instruments, persons of different temperaments will still see things a little differently. So in the final allowance for probable error which follows any calculation of delicate problems in astronomy, the astronomer takes his personal equation into consideration. But apart from this element of uncertainty, it is still not all a plain case of making and recording observations in seeking a parallax. Account must be taken of refraction, which causes an apparent deviation from the real position of an object; of the motion of the earth through space, which does not stop for the making of any observations, however important; of nutation, which is a wobbling of the earth on its axis; of precession, which is another motion of the earth; and of still other disturbing motions.

The determination of the distance of the stars is still of great importance to astronomers. However, indirect methods of determination are now being pursued more actively than the direct

method of determination. The indirect methods involve using certain characteristics of the star to give an estimate of its true luminosity. The spectrum of a star, for example, provides a good measure of the surface brightness. The relative intensities of certain lines can indicate the pressures within the star's atmosphere and enable the astronomer to estimate whether it is a large bright star or a small faint one. Or, in the special case of certain types of variable stars, the period of the light variation provides a measure of the star's actual luminosity. In either case, when we know the star's intrinsic brightness, and also the star's apparent brightness, we can then simply calculate how far the star must be from us in order to appear as bright as it does in the sky.

As previously noted, the brightest stars tend to be the nearest ones to us, but there are many exceptions to this rule. It is more accurate to say that, of the naked-eye stars, the faintest ones are farthest away. Alpha Centauri, the nearest star of all, is a brilliant first-magnitude star a little more than four light years away. Procyon, another first-magnitude star, is something less than ten light years away. Sirius, the brightest star in the heavens, is also comparatively near, its light journey through space being a little more than eight years (8.6). On the other hand, the second nearest star, usually called Barnard's star, is just a trifle brighter than the tenth magnitude (9.7) at a distance of about six light years.

Canopus, the star next brightest to Sirius, takes about 100 years to send its light to us. Arcturus, one of the brightest stars in the northern hemisphere, is 37 light years away. Spica is at a distance of 260 light years and brilliant Rigel at the record distance of 1,300 light years.

So, while it may be true that on the average a large proportion of the bright stars are nearer us than the faint ones, it does not necessarily follow that because a star is brighter than others it is, therefore, nearer, nor that any particular faint star is among those that are farthest away. Brightness is one of the criteria of nearness, but it is not always the determining one. The exceptions to this rule are due in a large part to the fact that stars themselves have intrinsic brightnesses ranging over an enormous number of magnitudes. There exist, in space, stars more than 10,000 times as bright as our sun. We also find stars 10,000 times fainter than our sun. These extremely faint stars are tremen-

dously abundant, but we can see them in our telescopes only to a relatively short distance.

Of the distances of the stars from each other we can have no conception until we know their distance from us. Some of those that appear to lie very close together may only lie in nearly the same line of sight and may really be separated from each other by many times their distance from us. There is reason to think, though, that many of the groups are not mere accidents of vision, but that most of the stars composing them have some close connection and are probably comparatively near each other. This is indicated sometimes through spectrum analysis and sometimes through telescopic observation. By the spectrum we learn that the stars in certain groups are all in nearly the same stage of development; and thus they seem to have had a common origin and may be comparatively near each other. This is the case with the stars of the Pleiades. By the telescope we learn that all the stars in certain groups are moving through space in the same direction; and thus they seem to have a common destination. Five of the seven principal stars forming the Big Dipper comprise one of these moving groups. The first star in the bowl of the dipper and the last star in the handle are not members. It turns out that our sun actually lies within the confines of this enormous moving cluster, although it is not a member. Sirius, Alpha Coronae Borealis, and Beta Aurigae are also members of the Ursa Major cluster.

The Pleiades and the Hyades are also examples of moving clusters. The motion of the latter group is rather unusual, in that the stars appear to converge toward a certain point in the sky. The effect is one of perspective. All of the parallel tracks in a railroad yard, for example, converge toward the same point on the horizon. In the same way, these stars, moving parallel to one another, converge to the same point, which is not far from the red star Betelgeuse in Orion. Many of the bright stars in the constellations of Scorpius and Centaurus belong to the same moving cluster. The beautiful cluster that comprises the constellation Coma Berenices is a naked-eye group. So, also, is the faint naked-eye cluster Praesepe, in Cancer.

.

## XX

### Double Stars

A GREAT many of the stars that we see as one with the naked eye are, in reality, composed of two, sometimes of three, and in a few cases of four or more stars. The reason that we see them as a single star is, either that they are too close together to show any space between them at the enormous distance they are from us, or that one of them is too faint to be visible without a telescope. Castor is an example of the first case. The two principal stars composing it are of the second and third magnitudes, and either of them could be seen with the unaided eye if they were not so near together. Their nearness, of course, is only comparative. Hundreds of millions of miles divide them; but their distance from us is so great as to make many hundreds of millions of miles dwindle to a mere point. Incidentally, Castor is actually a multiple star, having, in addition to the principal pair, several fainter companions. Regulus is an example of the second case. It has a companion star which is too faint to be seen with the naked eye, and yet the distance between the two stars is no less than that between some other stars that the eye can easily separate because they can both be seen by the eye alone.

Sometimes we loosely call a star double when we can really see without a glass the two stars composing it. Epsilon Lyræ, which we already know, is such a star. To many persons it seems to be a single star; but a good eye will discover that it is two. They are three minutes apart, and this is about the smallest distance that the eye can discern among the stars. A telescope, in turn, will resolve each of these stars into a close double, so that in reality Epsilon Lyræ is a double double system.

The star next to Aldebaran in the V-shaped figure of which it is a part is also spoken of as a double star, though its two

components are as much as six minutes apart and can ordinarily be seen with the unaided eye. There are many such double stars. But astronomers do not call a star double unless a telescope is required to separate the components of it, and even then they reject all stars that do not lie nearly ten times nearer together than the two principal stars of Epsilon Lyræ.

Sometimes two stars lie in such a position that they seem to us to be almost touching each other, yet are in no way connected and only happen to lie in the same line of vision, one being enormously farther away than the other. Their appearance as one is a mere matter of perspective. Such combinations of the light of two stars are called optical doubles, and are of no special importance as double stars in astronomical investigations after it has been fully established that they have no physical connection.

True double stars do not lie near each other by mere chance. They are held together by physical bonds that keep them forever revolving around each other, and their revolutions are what betray to us the relationship of the stars taking part in the motion. Sometimes it requires many years to determine whether there is a physical connection or not. For even though stars may be moving at the rate of many miles a second, their distance from us is so great that it takes a very long time for us to discover any motion at all. Hence, a star that requires several hundred years to complete its journey in its orbit would hardly reveal any perceptible motion in the lifetime of one man. A convenience it would indeed be to the modern astronomer if he could command the same consideration as the old historian Berosus says was given to the antediluvians when God prolonged their days to five or six hundred years in order to give them time to complete astronomical observations. The brevity of life, however, does not stay the inquiring mind. The work of investigation is passed on from hand to hand with almost as much continuity as if it were pursued by a single man.

Generally speaking, there is a real connection between two stars when they remain in the same relation to each other during a long series of years. For each star is in motion and has its own, or what we call its "proper," motion in the sky. If its companion has the same motion, and the two stars thus keep together, there is reason to suppose that they form part of the

same system. If they remain together and yet change their relative positions, we know that they are circling round each other and are undoubtedly of one system. But sometimes in their revolutions through their incredibly large orbits the two stars are enormously farther apart at one point than at another and may seem to be receding from each other. In such cases it requires long years of observation to determine whether each is going its own way and they will continue to recede, or whether they are bound together by gravitation and will in time approach each other again.

Sirius and its companion are entirely inseparable at one point in their circuit and at another can be separated by a large telescope. But as their period of revolution is only about fifty years, their relation was soon established after the smaller star was once seen during one of its periods of greatest distance from the principal component of the system.

Castor and its distant companion, on the other hand, can always be seen separately by the aid of the telescope. But it was only after they had been known as a double star for more than a hundred years that any physical association was suspected between them. It came about then by the accidental discovery that while once the prolongation of the line joining Castor and Pollux would run through Castor's companion, it was no longer so; but that the companion had moved quite out of line with Castor and Pollux, showing that the two stars must be in revolution. The time necessary for a revolution of a companion of Castor around its primary is slightly under 400 years.

Double stars that are not mere optical doubles, but are proved to have some physical connection, and, hence to revolve around some common center, are known as binary stars, or binary systems, and the stars composing such a double star are spoken of as its components. Those that can be separated with the telescope are called visual, or sometimes telescopic, binaries. But there is a class of binaries that cannot be separated even with the strongest telescope. Their double nature is made known to us by means of the spectroscope, and they are called spectroscopic binaries. Their discovery is one of the most interesting results of the use of the spectroscope in astronomy.

The displacement of certain lines in the spectrum of a star enables astronomers to determine the swiftness and direction of

its motion. In the case of a double star the components revolve around each other, alternately approaching and receding from us. The lines of their spectra indicate these motions and betray the fact that they are circling around in an orbit, and are, hence, double stars, since if they are revolving they must be revolving around something. When the two stars are of the same class spectroscopically, they will each have the same kind of spectrum and the lines of their spectra will alternately appear double and single, according as the stars are side by side to our view or in the same line of sight from us. When the two stars composing the binary are so different in their stage of development that one is nearly dark, the lines of the spectrum of the bright component will simply oscillate back and forth, for, of course, a star that does not radiate light has no spectrum. Mizar is an example of the first class and was the first star to be identified as of that class, this occurring in 1889. Mizar was also the first star discovered to be a double star. Spica is an example of the second class, since its companion star is almost non-luminous and signals its presence only by the wavering lines of Spica's spectrum. Capella is of a class between the two, where the spectra of both components can be seen, though they may be of slightly different spectral types.

Up to the middle of the seventeenth century such a thing as a double star was unknown. In 1610 Galileo first used his telescope, by which he discovered the moons of Jupiter. For forty years after that the heavens were scanned with various telescopes more or less rude in construction, but not once did it occur that a star which looked single to the naked eye became known to be double as viewed through a telescope. Finally, in 1650, Mizar was discovered to be composed of two stars. The idea once entertained, double stars became an object of special quest, and many others were discovered in quick succession, among them Alpha Centauri and the famous Theta Orionis, the star in the nebula in the sword of Orion, which was discovered as early as 1689 to be not only double, but quadruple. But no one dreamed that double stars were anything more than optical doubles. It was supposed that by mere chance both stars lay in the same line of vision from us, against the same background of black space, but one of them necessarily much farther away than the other.

More than a hundred years after the first double star was



discovered, and so recently as about the time of our Revolutionary War, a systematic search for double stars was begun. In less than two years thirty-three stars were added to the number already known. The astronomer Herschel then set about making a special study of double stars, in the hope that he might be able to learn something about their distance. The idea that one was far behind the other in our line of sight at least differentiated them as to distance and suggested the thought that it might be possible to discover some shifting of the nearer star in its relation to the farther one. The unexpected happened again, and a new scientific truth was discovered almost, as it seemed, by accident. The stars were revolving around each other! Another of nature's secret doors had been burst open by the continuous pressure upon it of zealous, untiring workers while chiselling away one fastening after another.

By 1802 it was proved beyond a doubt that many of the so-called double stars were truly double—twin suns swinging around a common centre of gravity: millions of miles apart, no doubt; but held by that mysterious force that keeps our own system together, in bonds that can never be loosened.

When Sir Isaac Newton termed that force "universal gravitation," stressing the fact that the force held everywhere in the universe, he was indeed far-seeing. These double stars proved beyond all question the universal character of the force that causes a stone to fall to earth, the moon to revolve in its orbit, or the planets to circle the sun. Here was implication that many other systems like ours could exist in the universe: a vast central orb holding smaller ones by invisible chains and swinging them around as a boy sometimes swings a pot of burning coals suspended upon a string.

Sometimes two fiery suns of almost equal size are circling around each other. They may each be the center of a planetary system and so far apart that they show as mere stars to each other. That they are so far apart, we know in some cases; and we even know that in some cases dark bodies are revolving around them. We may more appropriately refer to these dark bodies as unseen companions, because in most cases the bodies are not completely dark. They are only relatively faint stars which might actually be seen in a large telescope or photographed were it not for the presence of the bright companion.

Why may it not be that many others are thus far apart and accompanied by dark bodies which we have as yet no means of discovering? Scientists have even suggested that our sun may have a far-distant companion. However, the evidence is decreasing for the reality of a companion sun, but the possibility cannot yet be completely ruled out.

As soon as the fact that there was a physical connection between the components of some of the double stars was established, many workers began to devote themselves to the observation of such stars, sorting out and discarding the mere optical doubles, watching and recording the movements of those that proved to be real binary systems, and discovering new ones all the time. For more than a century the work was carried on by means of the telescope, and more and more pairs came into view as larger and larger telescopes came into use.

Astronomers have estimated that almost one-third of all the stars in the sky are binaries. The high incidence of binary systems clearly indicates that the evolutionary process leading to the formation of the double star is in no sense unusual. The indication is that a single star, originally rapidly rotating, may under certain circumstances split up into a double star into which most of the rotational motion has gone into the orbital revolution of one star about the other. These findings, as well as the discovery that binaries, too close to be separated even with the strongest telescope, could be recognized either through the spectroscope or by the light variation of the eclipsing binary, greatly enlarged the field of investigation.

Nearly a hundred of the binary systems discovered have proved to be multiple systems composed of anywhere from three to seven stars. Castor is a well-known multiple system. Polaris is a triple star, Mizar a quadruple, Theta Orionis divides into four bright stars, and Epsilon Lyræ is composed of four. From the frequency with which this happens, it is hardly surprising to find almost any binary resolve itself on further observation into a system of several stars. Many binaries that can be seen with a telescope have been shown by the spectroscope to have both components double, thus making a quadruple system.

Many orbits have been computed, showing periods of revolution from less than two days to more than a thousand years in length. The far-distant visual couples swing slowly and majesti-

cally around their common center, while the close companions whirl with almost unthinkable rapidity through their allotted orbits, for the shorter the distance between two stars the greater gravitational force they exert upon each other and the more swiftly they compel each other to revolve.

The fastest revolving couples are, of course, spectroscopic binaries, many of which are so close together as to be almost in contact. In proportion to the number of each, the orbits of more spectroscopic binaries have been determined than of the visual systems; for telescopic doubles must be observed until it is shown that they either are or are not in revolution, while spectroscopic systems are discovered *because* they are in revolution.

The shortest periods among spectroscopic binaries are a few hours; the shortest periods among visual binaries are a few years. It is interesting to know that the longest period found that has reasonable claims to accuracy is that of a fourth-magnitude star in the region between Altair and the square of Pegasus. It is estimated to be nearly sixteen hundred years; but it will require several hundred years of observation to fully verify this estimate. The longest period that has been computed and fully authenticated is about four hundred years, though much longer periods are known to be approximately correct.

Multiple stars grade into more complex systems, like the Ursa Major cluster, the Scorpius-Centaurus group, the Pleiades and Hyades, or Praesepe. The delicate naked-eye cluster, Coma Berenices, is another. In the last chapter we saw that such groups are physically connected because the stars are moving parallel to one another. The clusters hold their shape.

The discovery of binaries, or systems, of stars has a most important influence in the progress of astronomy. In establishing the universality of the law of gravitation it opens the way to knowledge of various sorts which would otherwise be unobtainable. When the distance of a pair of stars is known and the time that they require to revolve about each other, we secure data for determining the size of their orbit, their mass, their real light-giving power, and their rate of speed. In addition to this, if we learn from the spectroscope the chemical composition of the stars, it would seem as if their whole story is nearly within our grasp.

/

## XXI

### The Constellations

THERE is a general impression that knowing the stars means knowing the constellations; and with this idea in their minds many persons grow discouraged at the very outset in their endeavors to make some acquaintance with the stars. But the first step toward knowing the constellations is a familiarity with the individual stars. At first sight, most of the constellations do not much resemble the thing for which they are named. Many of the figures partially overlap each other. The tip of one of the horns of Taurus is in Auriga; Andromeda and Pegasus depend upon the same star to complete their outline; and in some other cases a star was traditionally assigned to two constellations, or families, so that it is difficult to tell of which one it is the step-child. Astronomers have, in questionable cases, arbitrarily decided to which star group a given star belongs so that the constellations do have officially recognized boundaries.

To enjoy the constellations one must be satisfied not to expect too much of them. The splendid ones that mark a certain section of the skies to us, as well as the small and beautiful ones that remind us by their presence that a certain season has arrived, yield their fullest pleasure when we take them as we do our human friends—for what they give us, without any severe exactions as to consistency or any desire to make them fit a preconceived model. The tracing of the various figures represented by the constellations was not the working out of an exact science, but it was an æsthetic occupation, and the imagination must be used to see, as the ancient star-gazers did, the heroes, kings, and queens, and birds and dragons fixed on the purple walls of the sky.

It is not necessary that one should be able to trace all the constellations in order to have a friendly and familiar acquaintance with all the important naked-eye stars. Still one can hardly be

said to know the aspect of the heavens until he knows the principal constellations—the ones that took shape in the minds of the shepherds, sailors, and other out-door people in the days when imagination fashioned them and when scientific order was of less import than it is to-day. These early forms include the twelve constellations of the zodiac, for instance, which mark the path of the sun and the moon and the planets across the heavens; the conspicuously beautiful constellations, like Orion, Lyra, Leo, Scorpius, and others; and the familiar ones forever circling around the north pole, and always visible to us, like Draco, Cassiopeia, and the two bears.

Moreover, however indifferent one may be to the science of astronomy, an intelligent person can hardly fail to feel a little uncomfortable when he has so large a body of natural beauty hanging over him in full view half of his life and yet has no familiarity with it. It was some feeling of this kind, no doubt, that led Thomas Carlyle to cry, "Why did not somebody teach me the constellations and make me at home in the starry heavens, which are always overhead, and which I don't half know to this day?"

Of the ancient constellations there are about forty-eight. They are all outlined by stars that can be seen with the naked eye and are as open to our vision as the clouds, the trees, and the hills. In them are contained all the brightest stars, and there is no conspicuous part of the heavens that is not covered by them. Once we are familiar with them, the whole sky becomes glorified to us beyond Achilles' shield, which Hephaistos fashioned and wrought thereon "the heavens, and the sea, and the unwearying sun, and the moon waxing to the full, and the signs every one wherewith the heavens are crowned, Pleiades, Hyades, and Orion's might, and the Bear that men also call the Wain, her that turneth in her place and watcheth Orion, and alone hath no part in the baths of Ocean."

Today, astronomers recognize 88 constellations. These comprise the 48 original constellations recognized by the ancients, with the exception of the Pleiades, which we now regard as being part of Taurus, and only an asterism. Also, astronomers have carved the original large star group, Argo Navis, the great ship in which Jason sailed to find the golden fleece, into four separate groups, the sail, the stern, the keel, and the compass. There have

been many new constellations added to the southern sky, regions invisible in the latitudes where the ancient astronomers made their observations. Also, the modern astronomers have straightened out the once irregularly curved boundaries until they now consist of arcs of circles.

Many of the modern constellations are of little interest or importance, having been composed to provide a scientific genealogy for the few inconspicuous stars occupying stray spots in the sky not covered by the original constellations. The ancient astronomers referred to these as the "unformed stars." But, for astronomical reference, it is often useful to have a map of the sky in which no blank spots occur. Hence, though the areas are generally poor in stars and often of little interest, they are necessary additions to the astronomer's system of mapping the sky.

Many of the newer constellations, especially those in the south, do not follow the ancient pattern. They are named for such astronomical or scientific instruments as the sextant, the octant, the telescope, the clock, the graving tool, the microscope and others. They are, in consequence, devoid of the romantic and historical interest that the ancient constellations commanded. Fortunately, the first astronomer to map the southern stars, Bayer, followed the ancient tradition and introduced groups of particular beauty and interest in the more brilliantly populated starry realms.

Fourteen of the principal constellations have already been described more or less fully in the introduction to the first-magnitude stars. With these as connecting links we shall proceed to identify and trace the other important constellations visible in the northern latitudes.

## XXII

### The Little Bear and Polaris

FIVE major constellations are always above the horizon in our latitudes: Ursa Minor, Ursa Major, Cassiopeia, Cepheus, and Draco. We can see them on any clear night from sunset to dawn. They are all in the north and revolve slowly in a small circle around the north pole once every twenty-four hours. In one sense the most important of these six constellations is Ursa Minor, or the Little Bear. The reason of its importance is that the north pole lies within its boundaries.

The north pole is nothing more than the point where the imaginary polar axis of the earth would touch the sky if it were extended that far. But since the earth revolves on its axis, the entire heavens seem to us to revolve around its extension; so the north pole is to our view the center of the universe and the most important point in the skies.

Of all the stars that can be seen by the naked eye the one nearest to the north pole is Polaris, or Alpha Ursæ Minoris. It is called the Pole Star, or the North Star, and is ordinarily regarded as marking the exact point of the north pole. It is, in truth, about one degree from the polar point and revolves around it each 24 hours in a little circle two degrees in diameter. A diameter of two degrees is nearly four times the apparent diameter of the moon, which is only a little more than half a degree. But the moon, by reason of its brilliancy, seems to cover more space on the sky than it really does, so we must make allowance for that fact in such a comparison.

To find the exact location of the pole, draw a line from Polaris to Mizar, the middle star in the handle of the Great Dipper. Mizar is on the other side of the pole from Polaris and almost exactly opposite; and the line will pass through the pole, which lies about one degree from Polaris.


Within the little circle that Polaris makes around the pole are many stars that are not visible to the naked eye. As many as eighty have been seen through the telescope, while as many as two hundred have been seen on the photographic plates.

Owing to what is known as the precession of the equinoxes, the pole does not always point to the same spot in the sky. It describes a little circle of its own once in every twenty-five thousand years, and as it journeys it confers the distinction of being the Pole Star upon several different stars. In about two hundred years from now it will approach our present North Star, Polaris, to within a little less than half a degree. The circle that Polaris will then make around the pole will have a diameter of about fifty-three minutes, which is slightly less than twice the diameter of the moon. After that, the pole will begin to move away from Polaris, and in about twelve thousand years will be within five degrees of the beautiful Vega, and she will be our Pole Star for the next three thousand years. Then the pole will slowly recede from Vega, and in another twelve thousand years Polaris will again become the Pole Star. So, in spite of all these changes, each generation during these thousands of years will have its Pole Star. For the change takes place so gradually that a single generation is not conscious of it. The change has been marked, however, over historical time. Some 2,000 years ago the "guard stars," Beta and Gamma Ursae Minoris, were nearer the celestial poles than was the star that we call Polaris. Hence it was customary for the early navigators to steer by the guard stars rather than by Alpha Ursae Minoris. Around 3,000 B.C., Alpha Draconis was the pole star.

Polaris is the brightest star in the constellation of Ursa Minor, or the Little Bear. Its Greek letter name is Alpha Ursæ Minoris. It was also called in former times Cynosura, meaning the end of the dog's tail. It is at the end of the tail of the Little Bear: and there is some reason to think that the Little Bear was once thought to be a dog instead of a bear. Our word "cynosure" gets its meaning from Polaris, which is the center of attraction in the heavens.

In the Little Bear, as in the Great Bear, the distinctive feature of the constellation is the Dipper. It is formed of seven stars and somewhat resembles the Great Dipper: but its handle is curved





THE LITTLE DIPPER

towards, rather than away from, the bowl, which gives it something more the appearance of a saucepan than of a dipper.

The only other stars of any importance in the constellation are the two outer stars of the bowl of the Dipper. The brighter one is a second-magnitude star, Beta Ursæ Minoris, sometimes called Kochab. The other is Gamma Ursæ Minoris, a third-magnitude star. They correspond in position to the pointers in the Great Dipper, and are sometimes called the "guardians of the pole."

## XXIII


# The Great Bear and the Seven Stars of The Dipper

UNLIKE the Little Bear, the Great Bear can be traced in the heavens with reasonable accuracy. It looks rather more like a rodent than a bear, and its long tail seems somewhat out of place. Still one can distinctly follow the outlines of a clumsily shaped four-legged animal, which, perhaps, is as appropriately called a bear as anything else. It is rather surprising, however, that many ancient peoples in widely separated continents have referred to this constellation as a bear.

The most conspicuous part of the constellation is the Great Dipper. The handle of the Dipper is the tail of the Bear, and the four stars forming the bowl lie in the hind part of the body. The tip of the Bear's nose is marked by a star lying about twenty-five degrees from the root of the tail, and a line from the nose through the two stars forming the top of the bowl of the Dipper marks the back of the Bear. Its right fore-foot is indicated by two stars about fifteen degrees south of its nose. The stars connecting this foot with the body show a prancing, double-jointed sort of a leg. Similar pairs of stars indicate the position of the other three feet, which can be appropriately connected with the body without much trouble, the whole forming an animal so huge that it might well give over its watch on Orion, since even so mighty a hunter as he would scarcely attack it.

Ursa Major is a beautiful constellation and would be remarkable even if its only distinction were that of including the seven bright stars of the dipper. These stars are well worth a closer acquaintance because of their many points of individual interest and beauty as well as their special interest as a group. With the exception of Orion, there is no other constellation with so many

bright stars in so close a group. Alioth, the third star from the end of the handle, is the brightest of them, and barely escapes being a first-magnitude star. All the others are of the second



THE SEVEN STARS OF THE BIG DIPPER

magnitude except Megrez, which lies at the point where the handle joins the bowl of the Dipper. This is a third-magnitude star.

Alioth is five and a half degrees from Megrez, or Delta, the faintest star of the seven, and is four and a half degrees from Zeta, or Mizar. The name Alioth means the "tail sheep."

Alkaid, the star at the end of the handle of the Dipper, is the third in order of brightness in the group, though no longer ago than 1847 it was said by Herschel to be the brightest. The difference in brilliancy between Alkaid and Alioth is hardly discernible by the ordinary eye. Alkaid is a white star and is approaching us at the rate of about seven miles a second. Its name means "the chief." The star is frequently called Benetnasch. It is seven degrees from Mizar, the star next to it in the handle of the Dipper, and twenty-six from Dubhe.

Dubhe, or Alpha Ursæ Majoris, is the pointer nearest the pole. The name means the "she-bear." Dubhe lies in the back of the Great Bear, ten degrees from the star that begins the tail of the Bear. It is the second in order of brightness in the constellation. Dubhe is yellowish in color and shows some similarity to Capella. It is a double star. Its smaller companion is about three magnitudes fainter than the larger star, being of the fifth magni-

tude. This makes a great difference in the brilliancy of the two stars, the brighter one being sixteen times brighter than the other. Dubhe is a little more than twenty-eight degrees from Polaris; ten degrees from Megrez, the other star at the top of the bowl of the Dipper; twenty-six degrees from Alcaid, the star at the end of the handle, and a trifle over five degrees from the other pointer. It is approaching us at the rate of five miles a second.

The other pointer is Merak, or Beta Ursæ Majoris. It is fifth in point of brightness in the constellation, its magnitude being 2.37, and is a very beautiful white star with a greenish tinge. It is approaching us at the rate of about seven miles a second.

From Phecda, or Gamma Ursæ Majoris, the other star at the bottom of the bowl, it is four and a half degrees to Megrez, or Delta Ursæ Majoris, the faintest star of the seven bright ones in the Dipper. Its magnitude is 3.30. Megrez is also whitish; like Phecda, it is slightly tinged with yellow. Its name means "the root of the tail," which describes its situation in the Great Bear. It is the star that joins the handle to the bowl in the Dipper. It is five and a half degrees from Alioth, the brightest of the seven, and is thirty-two and a quarter degrees from the pole.

The most variously interesting of all the stars of the Great Dipper is Mizar, or Zeta Ursæ Majoris, the star next to the one at the end of the handle. It is fourth in order of brightness among the seven, its magnitude being 2.05, almost an exact second-magnitude star. The light from Mizar takes about 80 years to reach the earth.

Mizar can always be known by its having a tiny companion so near it as to be sometimes not easy to see. Its name, Alcor, means "the near one." It was formerly known as "the test," because it was supposed to test the eyesight to see it. It is really not so difficult to see, however, for it is only a fourth-magnitude star and is separated from Mizar a distance equal to almost one-third of the apparent diameter of the moon. The greater light of Mizar is what makes it seem so obscure. When Alcor is once known, it never fails to attract, for it is a peculiarly beautiful little star, very white and very piercing in its brightness. We have already noted that all of the Dipper stars with the exception of Dubhe and Alcaid are members of a giant moving cluster. Alcor

is also a member of this group. Thus Mizar and Alcor are not too distant from one another.

But Mizar has other stars nearer it than Alcor. It is the center of its own great system of stars. It has the distinction of being the first double star discovered by the telescope, a discovery made as long ago as 1650. When the spectroscope came into use, Mizar was again the first to open up a new line of discovery, being the first star discovered by the spectroscope to be a binary. The brighter of the two stars into which it was separated by the telescope was shown by the spectroscope to be also double. So the star as we see it is really three stars all united into one system. The pair of stars that require a spectroscope to separate them revolve around each other in a little more than twenty days and radiate about forty times as much light as the sun.

In England the Dipper is often called the Plough and sometimes Charles's Wain, or simply the Wagon. As noted earlier, all of its stars except Dubhe and Alkaid are moving through space in the same direction and at about the same rate of speed, which indicates that they may all be of one system, or, at least, of a common origin, and thus be associated in reality as well as to our view. Since Dubhe and Alkaid do not belong to the cluster and are moving through space in a very different direction, the dipper will become somewhat misshapen in the course of centuries. In about 50,000 years it will appear more like a lopsided steamer chair.

.

## XXIV


### Cassiopeia, Cepheus, and the Dragon

CASSIOPEIA lies on the other side of the pole from Ursa Major and almost directly opposite the Dipper. Its chief naked-eye stars, seven in number, form a figure somewhat resembling a chair in shape. They are nearly as conspicuous among the circumpolar stars as the Dipper and are about the same distance from the pole.

On account of the faintness of the little star Kappa ( $\kappa$ ), on the front edge of the seat of the chair, some persons leave it out of consideration and fancy the remaining stars outline a somewhat distorted "W."

Gamma ( $\gamma$ ) Cassiopeiæ is at present the brightest star in the constellation, and is of the second magnitude. Its light varies irregularly, and for a short time it once rivalled Deneb in brightness. Alpha is a shade less bright, and Beta is also of the second magnitude. Delta ( $\delta$ ), the first star in the back of the chair, is close to the third magnitude in brightness. Epsilon ( $\epsilon$ ), at the top of the back, is of the third magnitude; and Kappa, already referred to as the little star on the front edge of the seat, is a fourth magnitude star. Eta ( $\eta$ ), half-way between Alpha and Gamma, is an interesting binary, or double star. The two stars composing it complete a revolution around each other every five hundred years, and are comparatively near to us, being only about eighteen light years away.

Only two stars in Cassiopeia have individual names. Alpha is sometimes known as Schedar, and Beta was formerly called Caph. Polaris is just about half-way between Caph and Megrez, the star that joins the handle to the bowl of the Dipper, and is on the same side of the pole with Caph.



CASSIOPEIA

## CEPHEUS

Next to Cassiopeia is Cepheus, the king, her husband. He stands with one foot on Polaris and his head reaching to Andromeda. The constellation has only one second-magnitude star, Alpha, which was formerly called Alderamin, meaning "the arm." This star and Beta, a rather bright third-magnitude star lying eight degrees nearer the pole than Alpha, point almost as directly to Polaris as the well-known pointers. They are a little nearer the pole than the pointers and are about three degrees farther apart.

When Cassiopeia is east of the pole Cepheus is just above her. When she is above the pole, he is west of her; and when she is below it, he is just east of her.

## THE DRAGON

The only other important constellation among those around the north pole is Draco, or the Dragon. It is supposed to represent, and does somewhat suggest, a great dragon that winds in

and out among the circumpolar stars in graceful serpent-like convolutions. It is the Dragon on which the giant Hercules rests one foot. Its head is marked by a four-sided figure a short distance northwest of Vega, containing two bright and two faint stars. Since the head of Draco never sets, it is interesting to watch it when Vega is below the horizon, and one sometimes has a feeling that by a good big stretch on the tip of his toes he could see the bright Vega just below the shining eyes of the Dragon. The rest of the constellation is outlined by a number of faint stars. It runs a short distance towards Cygnus, and then with some well-marked undulations it runs up towards Cepheus until it is between Cepheus and the Little Dipper. There it turns, makes a sweep around the Little Dipper, and, running between it and Ursa Major, ends just above the bowl of the Great Dipper.

The two brightest stars in Draco are the ones in the head. The brightest is the one nearest Vega, which is Gamma ( $\gamma$ ) Draconis, sometimes called Etamin. The other is Beta ( $\beta$ ), with the individual name of Alwaid. Gamma is a second-magnitude star, and Beta is of the third magnitude. Gamma is a beautiful orange-colored star and is about 80 light years away from the earth.

A small star near the tail of the Dragon, lying half-way between Mizar, of the Great Dipper, and the lower star in the bowl of the Little Dipper, is noted as having been the Pole Star four thousand years ago. Its name is Thuban, and it is the Alpha of the constellation, though Draco now contains a number of other stars that are brighter.



## XXV

### From the Hunting-Dogs to the Arrow

#### CANES VENATICI

**J**UST under the handle of the Great Dipper is a collection of faint stars called Canes Venatici, or the Hunting-Dogs. The only naked-eye star of any importance in it is Alpha, commonly called Cor Caroli. It gets its name from a modern attempt to make a constellation in honor of King Charles II. of England. The name means "the heart of Charles." When the handle of the Great Dipper is turned towards the west, Cor Caroli hangs under it in a very conspicuous position and is a star that one would naturally notice and inquire about. It forms a triangle with Arcturus and Alkaid, or Benetnasch, the star at the end of the handle of the Dipper. It is one of the brightest of the third-magnitude stars. It is also a double star, with a companion of about the fifth magnitude; but the two are too close together to be separated by the naked eye.

#### BOÖTES

Canes Venatici is a sort of connecting-link between the strict circumpolar constellations and those that make a somewhat larger circle around the pole and are not visible all the time. Perhaps the easiest way to find them is to look for them one by one as they follow each other up from the eastern horizon. Canes Venatici is supposed to represent two hunting-dogs on the trail of the Great Bear, and they are naturally followed by the hunter, or the bear-driver, Boötes, which rises next. The constellation is represented by a kite-shaped figure that rises about eight o'clock early in March. One can find it easily by the great star Arcturus,

with which we are already familiar, and which lies in the knee of the hunter. It is a very pretty constellation and of some astronomical interest, but it is not remarkable to the ordinary view except for its one brilliant star.


## CORONA BOREALIS

Corona Borealis, or the Northern Crown, is a semicircle of stars that rises just east of the northern part of Boötes. Its chief star Alphecca (Alpha Coronæ) comes into view just an hour and a quarter later than Arcturus. It is of the second magnitude, though much fainter than the brightest second-magnitude stars. The constellation as a whole is a rather brilliant one, notwithstanding that it has no star brighter than the fourth magnitude except Alphecca. There are seven principal ones, and most of them are of the white, twinkling kind, so that the crown is full-jewelled. The entire constellation is only about fifteen degrees across.

## HERCULES

Next after Corona rises Hercules, one of the large constellations. It begins just south of the head of Draco and runs to within four degrees of the equator. The chief stars composing it delineate a giant upside down, as shown in the diagram on the opposite page.

This constellation represents Hercules kneeling and with one foot on the head of the dragon, Draco. Many authors have placed a club in his hand, but the concept of a kneeling giant carrying a club seems somewhat out of place. The reviser of this book (DHM 1963) has made a special study of the origin of this constellation. He believes that he has found the answer in diagrams and pictures of Hercules drawn on ancient pottery of Greece. Apparently the proper representation is that Hercules holds in his outstretched hand a bow. He is kneeling to gain stability. He has just shot the arrow, the constellation Sagitta, which is flying towards two birds, Cygnus and Aquila. The pottery fragment clearly showed the kneeling Hercules holding the bow, the flying arrow, and the Stymphalian birds as the



target. Many of the older sky maps represented Lyra also as a bird, a vulture. An alternative name for Lyra in these older maps was Vultur.

In the legends of Hercules we find that one of the tasks he was assigned during the performance of his twelve labors was the shooting of the birds that haunted the marsh Stymphalis. These birds were fierce, brazen-clawed and brazen-beaked. It seems entirely fitting that the constellation should depict the

famous giant performing one of his twelve great labors.

Hercules has no star as bright as the second magnitude, but has many of the third and fourth magnitudes, and has also several double stars. The constellation is directly north of Scorpius.

## LYRA AND CYGNUS

Lyra, or the Lyre, is the next constellation eastward from the northern part of Hercules. It contains the brilliant Vega, and its form has already been described. In addition to Vega, it has a famous star, Epsilon ( $\epsilon$ ); a variable star, Beta ( $\beta$ ), whose changing light results from the eclipses of two stars revolving around one another; and a rare nebula, known as the Ring Nebula, on account of its shape. Epsilon can be divided into two stars by a sharp eye, and to try to see them is an interesting exercise. Each of the two is again double. Vega and Epsilon, with Zeta ( $\zeta$ ), the star nearest to Vega in the parallelogram described on page 27, form an equilateral triangle each side of which is two degrees in length. This will enable one to find Epsilon.

According to legend, Lyra represents the invention of the musical instrument by the god, Mercury. He found a tortoise shell and strung strings across it, finding that the plucking of the strings gave out sweet music. One is not quite sure whether Lyra was supposed to represent the musical instrument itself or the tortoise whose shell became part of the instrument. Most of the early maps show it as a stringed instrument, however. But the representations consistently add wings and a bird's head to the instrument itself. This probably comes about because of confused mythology. We have already noted that Lyra sometimes has been referred to as Vultur, the vulture. Hence the bird's head may very well represent the head of the vulture. On the other hand, a badly drawn tortoise head might readily have been mistaken for the head of a bird in some artistic rendition of the constellation. So the beginnings of this beautiful group are lost in prehistory.

Cygnus, the Swan, is the constellation that holds Deneb. In its rising, Cygnus follows Lyra, and has been fully described in the chapter on Deneb.

## AQUILA, THE DOLPHIN, AND SAGITTA

In Aquila we recognize the bright Altair, which, with its two companions, already known to us, serves to indicate the constellation. Exactly eight degrees below Altair when it is on the meridian is a conspicuously variable star (Eta Aquilæ), which changes from the third to the fifth magnitude every seven days. Its changes are well-marked to the unassisted eye and make it an interesting object to follow from night to night. It is best seen in the early autumn, when Aquila is on the meridian during the first part of the evening. The two companions of Altair are Beta and Gamma Aquilæ. Beta is south of Altair, and Gamma is north of it. The line joining the three stars is just five degrees in length. Aquila is supposed to represent an eagle flying towards the east and across the Milky Way.

The Dolphin, or Job's Coffin, in the shape of a diamond, is just east of Aquila and a little farther north than Altair. Three of the stars composing the diamond are a little variable; one of them is a double star, and another is a spectroscopic binary, which is also suspected of being a triple star.

Near the Dolphin is an equally miniature constellation, Equuleus, the small horse. This constellation lies in the area of the sky devoted to aquatic creatures. The reviser of this book, therefore, prefers to regard Equuleus as a Sea Horse rather than an ordinary horse. This makes it an appropriate companion to the nearby Dolphin.

Sagitta, or the Arrow, lies in the Milky Way, north of Aquila and south of Cygnus. It is composed of five stars so arranged as to suggest an arrow pointing towards the east. Though a very small constellation, it is pretty and easily found. As previously noted, Sagitta naturally fills the position of the arrow shot from the bow of Hercules.

## XXVI

### From the Winged Horse to the Bull

THE next constellation of any importance that we may expect to see in the east is Pegasus. Nearly three hours after Altair is in view, the principal stars of Pegasus are seen coming up over the horizon a little north of east. It is a large constellation composed in the main of small stars. The chief figure consists of a large square called "the square of Pegasus." Each side is nearly eighteen degrees long, and the four stars composing it are sufficiently bright to make the square a significant figure in the sky.

On the northeastern corner of the square is a small triangle which serves as an identifying mark. The star on this corner of the square is called Scheat. It is Beta Pegasi and the third star in order of brightness in the constellation. Alpha Pegasi (sometimes called Markab, the "Saddle") is on the southeastern corner of the square. Alpha is slightly brighter than Beta. Algenib, or Gamma Pegasi, marks the other lower or southwestern corner. This star is somewhat less bright than Beta. The star in the fourth corner of the square, which is the brightest of them all, really belongs to Andromeda, though, in effect, two constellations share the star. It was formerly Alpha Andromedæ and also Delta Pegasi: but it is now not included in Pegasus, although it is clearly necessary to complete the asterism called the Square of Pegasus. Its individual name is Alpheratz. The brightest star belonging to the Pegasus family is Enif, or Epsilon Pegasi. It lies about ten degrees east of the little diamond of the Dolphin.

Pegasus is supposed to represent the winged horse, the favorite of the Muses. But as to its position in the sky authorities differ. Some depict the horse as rising upside down, with his fore-feet pawing the air up towards Cygnus and the top of his head resting on the urn of Aquarius. But if one adopts that position, Pegasus never entirely rights himself in his journey across the skies, as

some of the figures of other constellations do. In this representation, ever on his back, he goes pawing across the heavens and down "into the depths of the ocean"—symbolic, perhaps, of the condition ascribed to those who ride him.

However, there is an entirely different picture, one more consistent with the mythology of the group. In this representation the square no longer outlines the body but the wings. The entire figure is reversed, top for bottom. The important effect of this inversion is that the front feet of the winged horse now rest directly on Aquarius. The legend is that the horse struck the rocks with his forefeet, causing a fountain to flow from them, the fountain of Hippocrene. The reviser of this book prefers this second representation.


## ANDROMEDA

The constellation Andromeda is supposed to represent a beautiful lady chained to the rocks. She was the daughter of Queen Cassiopeia and King Cepheus, who are near her in the skies and have been described, and she was brought to this situation because her mother boasted too much of her beauty.

The constellation is more than thirty degrees in length and is outlined by a row of four bright stars, beginning with the one in the corner of the square of Pegasus and running northeast across the heavens in a slightly curved and easily traced line under Cassiopeia. Andromeda and the square of Pegasus are shown in the illustration, together with the "segment of Perseus," which lies next to Andromeda.

The main stars in the two constellations are thought to suggest an immense dipper; but for that matter one can find a whole tinshop of dippers in the sky if his mind runs that way and he is willing to admit a variety of styles in this useful utensil.

Andromeda rises cradled on the rock to which she is bound, her head lifted gently into view by the great square of Pegasus, on which it rests, and her feet, very properly, coming up last. But she sets in a most undignified fashion, drawn down head first by the flying Pegasus, her feet waving above the horizon after the rest of her is out of sight. The four chief stars forming the



ANDROMEDA, THE SQUARE OF PEGASUS, AND THE SEGMENT  
OF PERSEUS



curved line are Alpheratz, or Alpha Andromedæ, which marks the head; Delta, lying in the shoulder, from which an arm is outstretched towards the north; Beta, or Mirach, marking the girdle; and Gamma, or Almach, indicating the foot. They are all second-magnitude stars except Delta, which is of the third magnitude. Almach is a noted triple star which can be separated by the telescope. The components are orange, emerald, and blue.

A short distance north of Beta is the famous Andromeda Nebula, the brightest nebula in the entire sky. To the naked eye it appears as a fuzzy star. A binocular, however, or a small telescope will show it as an oval patch of haze. This object, M31 and its faint companion M32, are galaxies, Milky Way systems like our own. The larger contains perhaps a hundred billion stars, revolving in a sort of stellar pinwheel. Light from the Andromeda Nebula takes about one and a half million years to reach us. Large telescopes break the structure up into separate stars, though the hazy background of light comes from the billions of stars too faint to be seen individually.

The Andromeda Nebula has one other distinction. Since you can see it with the unaided eye, it is undoubtedly the most distant object that anyone can see without telescopic aid. The eye, however, even with the help of the largest telescope, does not show the true spiral structure and individual details. It takes photography to reveal the beauties of this nebula, which is the nearest example of billions of other spiral galaxies, individual clouds of stars that lie, here and there, like oases in the empty universe of space. In the southern sky appear two nearer galaxies, irregular in shape, the large and the small Magellanic Clouds. These two galaxies are about 100,000 light years distant.

## PERSEUS

The hero that rescued Andromeda from her woeful plight was Perseus, and he also is enshrined in a constellation that adorns the sky in the region a little north of Andromeda. Perseus lies in a very brilliant part of the Milky Way and is a beautiful constellation. The principal figure in it is a curved line of stars called the "segment of Perseus," running from Cassiopeia towards

Capella. Near the center of the segment is Mirfak, or Alpha Persei, a bright second-magnitude star, the brightest in the constellation.

The most noted star in Perseus is Beta, or Algol. It lies a little southeast of Mirfak and forms a triangle with it and Almach, the star in the foot of Andromeda. Algol varies in brightness from second to third magnitude once in about every three days (2 days, 20 hours, 48 minutes, and 55 seconds).

The name Algol signifies "the Ghoul." Indeed, the original representation of Perseus depicts him as holding in his hand the severed head of the Gorgon Medusa, the sight of which was supposed to turn a person into stone. It is doubtful indeed that the star received its name from the peculiar variation. The ancients were probably unaware of the "winking" that seemed to lend to the star a sort of demoniac quality.

Since Algol varies at such short intervals and so markedly, it has long been an object of interest even to the naked-eye observer. The mystery of its changes has been made clear by the spectroscope, and it is now known to be a spectroscopic binary with an invisible companion that eclipses it about once in three days. The eclipse lasts about eighteen minutes, and for that length of time it is a faint third-magnitude star. Then it regains its brightness and remains a second-magnitude star until the next eclipse.

Algol is interesting to observers in the latitude of New York City as being the only star as bright as the second magnitude that comes very near the zenith, or the point directly over our heads, on its course across the heavens. When on the meridian it is only about one-tenth of a degree from our zenith. Algol rises about an hour earlier than Capella. Perseus has two very beautiful star clusters which make a splendid display with the aid of ordinary opera-glasses.

## AURIGA

Auriga rises next after Perseus in the northeast. It is the constellation that contains the charming Capella, which is, of course, its brightest and most important star. Its five brightest

stars outline a five-sided figure that somewhat resembles a shield. Aside from Capella, the constellation has only one star as bright as the second magnitude. This is Beta Aurigæ, or Menkalinan. The other three stars of the pentagon are of the third magnitude.

The mythological significance of Auriga is somewhat confused. Many of the atlases depict Auriga as a goatherd holding a small she-goat in his arms. Indeed, the term "Capella" does mean the "goat star." There is the possibility, however, that the goat star received its designation because its appearance in the sky presaged the rising of the constellation of Capricornus, the Sea Goat. Capricornus was supposed to bring torrential rains and hence the recognition of its rising was important to the ancients.

On the other hand, there is an entirely different potential representation of Auriga. Many authorities have depicted Auriga as the Charioteer. The question is, then, who is riding in this chariot and why? Resolution of this enigma has been a particularly interesting problem to the reviser of this book. The answer seems to be relatively simple. The charioteer is none other than Neptune or Poseidon, god of the sea. Whenever the sea god rose from the waves and ventured forth on land he was supposed to ride in a chariot drawn by sea horses. Or, in other representations, this chariot was a sort of sea chariot drawn by sea horses whose reins Neptune held in his hands. In the vicinity of ancient Rome, many fountains or mosaics depict Neptune rising from the sea, riding in the chariot and driving the sea horses before him.

How appropriate it is that Neptune should be in this particular place in the sky. For indeed it was Neptune who formed the base of the entire legend linking the constellations of Cassiopeia, Andromeda (daughter of Cassiopeia), Perseus, and Cetus. It was Cassiopeia who offended the sea nymphs by boasting that she was fairer than they. Thus she was banished to the sky, to swing around the pole, half time head downward. And, as additional punishment, Neptune chained her daughter to a rock and sent the Sea Beast, Cetus, to ravage the coast and devour the maiden. It was Perseus, riding in on Pegasus the winged horse, who rescued the maiden. The lost meaning of the constellation of Auriga is thus reconstructed. The five-sided figure may

well be the chariot on which Neptune is riding, surveying the havoc his orders produced.

## TRIANGULUM

Just below, or east, of Andromeda is a pretty little constellation called Triangulum, or the Triangle. There are only five stars in it, and the brightest one is only of the third magnitude. Two others are of the fourth, and two of the fifth magnitude. It rises about half an hour after Andromeda.

## ARIES

Next east of Triangulum is Aries, a small but important constellation. Its importance consists in its being one of the constellations of the Zodiac, which mark the path of the sun. The sun passes through it from April 15th to May 13th; so that six months later it is exactly opposite the sun in the skies and rises just as the sun sets. It is the first sign of the Zodiac; but owing to the precession of the equinoxes the signs of the Zodiac do not now correspond with the constellations of the same name. From the middle of March to the middle of April, when the sun is in the sign of Aries, it is in the constellation Pisces; and when it is really in the constellation Aries, it is said to be in the sign Taurus.

The chief stars of Aries form a triangle in size about like Triangulum, and the two constellations are sometimes mistaken for each other by observers not entirely familiar with these constellations. It should be remembered that Triangulum rises first and is about half-way between Aries and Andromeda, and that the apexes of the two triangles point in different directions.

The chief star of Aries is Hamal. It rises about three-quarters of an hour after Algol and about fifteen minutes before Capella. It is the brightest star in the constellation and is of the second magnitude. The next brightest is Beta Arietis, or Sheratan. It is about one-half a magnitude less bright than Hamal and is about three degrees below, or east of, it in the sky. The two stars mark the base of the two horns of Aries, or the Ram.

## TAURUS

Taurus is the splendid constellation in which Aldebaran lies, and can always be recognized by that bright star. It is the third constellation and second sign of the Zodiac. The sun passes through it from May 13th to June 18th, so it may be seen rising as the sun sets six months later—from the middle of November to the middle of December.

Taurus contains the cluster of the Hyades, which are supposed to rule over wet weather and are sometimes referred to in poetry as the "moist Hyades." Aldebaran, as we have seen, is one of this cluster. The cluster outlines the face of the Bull. Above it are the Pleiades, which lie in the Bull's shoulder. The tips of the horns are indicated by two stars near Auriga. One of these stars is shared by both constellations to complete the legendary outlines. However, it officially belongs to Auriga.

The Bull rises backward, with its head lowered under the threatening arm of Orion, and it backs all the way across the heavens, going down without having made the terrible onset that it all the time promises.

## XXVII


### The Mystery of the Pleiades

OF all the groups of stars in the heavens none has excited so universal and so romantic an interest as the Pleiades. The magic of their quivering, misty light has always made a strong appeal to men of imagination. Minstrels and poets of the early days sang of their bewitchment and beauty, and many of the great poets, from Homer and the author of Job down to Tennyson and the men of our own day, have had their fancy enlivened by them, and in one form or another have celebrated their sweetness and mystery and charm. Men of fancy have compared the Pleiades to a swarm of fire-flies, to bees, to a rosette of diamonds, and to shining dew-drops, while less ecstatic minds have compared them to a hen and chickens and the seven virgins. Even people who do not know them by sight and have not felt "the sweet influences of Pleiades," vaguely recall some story about a lost Pleiad that stirs an emotion suggesting something romantic and sad.

The Pleiades form, in truth, a delightful group of twinkling, unfathomable stars, singularly fascinating and singularly persistent in their brilliancy. Of the six that can usually be seen with the naked eye, all but one are of the fourth magnitude and are not much brighter than the tiny Alcor, Mizar's companion. Yet often when the moon is full and many of even the third-magnitude stars are practically extinguished by its greater brilliancy, the Pleiades hang in the sky like a small luminous cloud distinctly visible and commanding admiration. When the moon is dark, they are hardly less conspicuous than a first-magnitude star, and one sometimes fancies what a splendid great star they would seem to us if they would only huddle a little closer together so that we could see them as one.

Still, even on the surface of the sky just as we see them, they

are not so close together as we often regard them. The full moon, which to casual observation seems to cover an area so much greater than they, can move about among them, most of the time covering only one visible star at a time. This occurs when the Pleiades are occulted by the moon, a phenomenon that is sometimes seen, since the Pleiades lie within four degrees of the center of the Zodiac, in which, as we know, lies the path of the moon. The whole constellation covers a space of a little more than three square degrees, while the apparent diameter of the moon is but little more than half a degree. It is hard to believe that four or five moons together would be required to cover the little dipper of the Pleiades from our view entirely, but it is true. The moon appears much larger to us because of its brilliancy. Its size as



THE PLEIADES

compared with other objects in the heavens is very deceptive.

The principal stars of the Pleiades form an outline resembling a tiny short-handled dipper. I have more than once been asked if this was the Little Dipper. The Pleiades, of course, lie in an entirely different place from the true Little Dipper, the one that has Polaris at the end of its handle. The Pleiades lie in the constellation Taurus. They rise just an hour earlier than Aldebaran and are visible at the same seasons as that bright star. We see

their shining light during the first cold evenings of autumn. On Hallowe'en they are about half-way up from the horizon between nine and ten o'clock. In the winter, the Pleiades are high in the sky during the early evening, tending more and more toward the west as the season advances. In March they set between nine and ten o'clock, and in April they disappear from the evening skies.

Any one can see the six principal stars of the Pleiades; many can see seven, and sharp eyes can distinguish anywhere from eight to eleven. There are so many stars in the group that are just beyond the limit of easy visibility that to any eye the number varies according to the clearness of the atmosphere and other physical causes. Even from moment to moment they twinkle in and out of our vision. It was no doubt this characteristic that suggested the simile of a "swarm of fire-flies." There are, in fact, more than three thousand stars in this small group. They cannot all be seen with a telescope, but they have recorded the fact of their existence on the photographic plates.

Photography, too, has disclosed the extent and complexity of a vast body of nebulous matter—"a cosmical fog," it has been called—which hangs around the Pleiades like a great veil, gathering itself into wisps here and there and in some instances making something like festoons from one star to another. Slight traces of this great cloud of nebula were first seen visually through the telescope, but only within the last few years have the extent and significance of the enveloping cloud been learned.

The chief stars of the Pleiades group have the same general motion in the heavens, and more than forty of the fainter ones have been observed and are found to be moving along with the system.

The Pleiades appear to be a great system of stars, perhaps evolving from the tremendous mass of nebula in which it is engulfed. In comparison with Alcyone and other Pleiades our sun is small indeed, and at the same distance would be visible only with a good telescope. The distance of the Pleiades from us is about 400 light years.

In keeping with the magnificent scale on which the whole system is constructed is the distance of the individual stars in it from each other. The diameter of the cluster is of the order of thirty light years. Yet so distant are they from us that with this



vast space between the individuals the group shows to us only the tiny cluster of twinkling stars, small enough barely to cover the jewels in Orion's belt.

We may know more about the Pleiades as time goes on, but they will never be any nearer to us or any brighter to our view. They may indeed in future ages be even less bright, for every second of our lives the distance between us and them is increased by many miles. Our sun, with us, his children, is travelling through space away from the Pleiades at the rate of about six miles a second, while they are headed towards another part of the universe and travelling at no less a rate. And yet, in seeming contradiction to this, under an old Hindu conception of the material universe, Alcyone, the brightest star in this group, was supposed to be the center around which our sun with all its planets and other suns with their planets were revolving in immense orbits. This idea is not correct, even though it reappears perennially, supported not by astronomers or scientists but by speculative philosophers expounding the metaphysical thought of the East.

The nine brightest stars of the Pleiades bear the names of Atlas and his seven daughters and their mother, the nymph Pleione. Atlas is the star at the end of the handle of the dipper formed by the stars. Pleione, the mother, which lies just above him, may be the lost Pleiad, since it is difficult for some persons to see her. Alcyone, the brightest of them all, is of the third magnitude, and lies at the joining of the bowl and the handle of the dipper. Electra, the second star in point of brightness, lies in the bottom of the dipper and is of the fourth magnitude. It is but a shade brighter than Atlas, and the ordinary eye can see no difference between the two. Maia, which lies about the middle of the top rim of the bowl; Merope, in the back corner of the bowl, and Taygeta, at the front edge of the brim, are all fourth-magnitude stars. Celæno, a little fainter, is about half-way between Electra and Taygeta; and Asterope, the faintest of her family, lies a short distance above Taygeta. Asterope is of the sixth magnitude, and very few persons can see it without the aid of a glass.

From the persistence of the story of the "lost" Pleiad and the almost universal reference to them as "the seven stars," it has been suggested that one of the fainter ones may formerly have

been brighter, but it is not certain which one. Electra is sometimes mentioned as having veiled her face on account of the destruction of Ilium, and sometimes as having been carried off by Canopus. If she is the lost one, it is in no physical sense, for she still shines the second in brightness of them all. The like is true of Merope, often reputed to be the lost one because she married the mortal Sisyphus, but she is still among the six brightest Pleiades. Celæno has been reported to have been struck by lightning and thus lost, fading to the seventh magnitude and therefore difficult to see with the naked eye. Pleione, the mother star and of the seventh magnitude, is the one commonly assigned the role of the lost Pleiad.

## XXVIII

### From the Twins Back to the Hunter

GEMINI is the constellation of the Twins, already described. It rises about the same time as Taurus and is next north of it. Gemini is the fourth constellation and third sign of the Zodiac. The sun passes through it from June 18th to July 18th. Hence it can best be seen from October to May.

#### CANIS MINOR

Canis Minor lies south of Gemini and is already known to us. It is the constellation containing Procyon; and though it has seven other stars, none of them is of any particular interest. It represents a very small dog on the back of a unicorn. The Unicorn is a constellation in itself, known as Monoceros, but it is insignificant and contains no star brighter than the fourth magnitude.


#### CANCER

This is a small constellation containing no star brighter than the fourth magnitude, and out of its fifteen naked-eye stars only four are as bright as that. It is the fourth sign and the fifth constellation of the Zodiac. After leaving Gemini, the sun passes through Cancer from July 18th to August 7th. The constellation includes most of the stars in the region between Pollux and Regulus. Its most interesting feature is Præsepe, a beautiful star cluster shaped somewhat like a beehive and lying about ten degrees south of Pollux. It is visible to the unaided eye and a particularly beautiful object in a small telescope or binocular.

## LEO

Leo the Lion is the constellation containing the first-magnitude star Regulus, with which we are already familiar. Leo is the fifth sign and the sixth constellation of the Zodiac, and the sun is travelling through it from August 7th to September 14th. It is best seen from January to June. It is supposed to represent a couchant lion and is one of the rather conspicuously displayed constellations.

It contains three stars of the second magnitude, Beta, Gamma, and Delta Leonis. Gamma is the brightest star in the blade of the Sickle. Beta, more frequently called Denebola, Delta, some-



LEO OR THE LION

times called Zosma, and Theta form a conspicuous triangle. Zosma lies in the back of the lion, at the beginning of the tail, and Denebola marks the tuft at the end of the tail. The triangle is visible in the northwest in the early summer for nearly two hours after the Sickle has set.

## COMA BERENICES

Just east of Leo is a small cluster of stars called Berenice's Hair. It is interesting mainly because of its delicate beauty.

This brings us back to Boötes in our circuit of the constellations south of those surrounding the pole and north of the equator. The next round will include the constellations coming within our view that lie either in part or wholly south of the equator.

## XXIX


# Constellations Wholly or in Part South of the Equator

## VIRGO

**S**OUTH of Coma Berenices and Boötes lies Virgo, the sixth sign and the seventh constellation of the Zodiac. It lies about half north and half south of the equator and is one of the largest of the constellations, but not one of the richest in stars. Virgo contains only about thirty-eight or forty naked-eye stars within its boundaries. One of these is Spica, already known to us. The next brightest is Gamma, a remarkable binary lying about ten degrees north of Spica. It has a period of one hundred and eighty-five years. It is of the third magnitude. All the others are of the third magnitude or less. The sun occupies Virgo for forty-three days, passing through it from September 14th to October 29th. Its stars being comparatively faint and much scattered, Virgo is difficult to trace. It is supposed to represent a goddess with wings, bearing a sheaf of wheat in one hand. Spica, as we have noted, represents the ear of wheat. The constellation occupies all the space from Leo to Libra on the west and east, and from Boötes on the north to Hydra on the south. Denebola, Spica, Arcturus and Cor Caroli form a large diamond, almost fifty degrees in length, called "the diamond of Virgo."

## LIBRA

Libra, or the Scales, is a small constellation lying wholly south of the equator, into which the sun passes when it leaves Virgo on October 29th. It occupies this constellation until November 21st.



LIBRA OR THE SCALES

Libra has no star brighter than the third magnitude. Its chief stars form a four-sided figure that is easily identified.

Libra is visible during the summer months crossing from southeast to southwest, where it disappears in November.


## SCORPIUS

Scorpius is known to us through its brightest star, Antares. It has in addition four stars of the second, seven of the third, and nine of the fourth magnitude. It is the ninth constellation and the eighth sign of the Zodiac, and is occupied by the sun from November 21st to December 16th. Scorpius is the most southerly of the constellations marking the path of the sun, and is farther south than any other constellation of which we have a complete view. But inasmuch as the sun does not pass through the most southerly part of Scorpius, it is not the constellation that con-

tains the point farthest south in the Zodiac. We can see it from about June 1st until late in October.

## SAGITTARIUS

Sagittarius is the ninth sign and tenth constellation of the Zodiac. In it lies the most southerly point of the Zodiac. The sun enters it on December 16th and reaches the most southerly point of its path five days later, December 21st. Sagittarius lies in the Milky Way east of Scorpius. It can be easily recognized by a little figure resembling a dipper turned upside down, an asterism known as the "Milk Dipper" because it lies in the Milky Way. It is very conspicuous during August and September.



SAGITTARIUS OR THE MILK DIPPER

Sagittarius represents a centaur in the act of shooting an arrow. The curved line of stars represents the bow; the one extending into the Milky Way and the one farthest east in the Dipper represent the arrow and the bent arm pulling the cord. Sagittarius has two stars of the second magnitude, nine of the third, nine of the fourth, and twenty-eight of the fifth magnitude. The

Milk Dipper is almost seventy degrees directly south of Vega and is almost as far south of the equator as Vega is north of it.

## OPHIUCHUS AND SERPENS

Ophiuchus and Serpens also lie on both sides of the equator. They form a sort of combination constellation that occupies the space between Hercules on the north and Scorpius on the south. Ophiuchus and Serpens contain no very bright stars. The combined figure represents a man bearing a serpent. The two legs of Ophiuchus run down to Scorpius, and the head bumps into the head of Hercules, which is on the southern end of that constellation. The serpent is more easily traced. Its head is just south of the Northern Crown, and thence it winds down alongside the head of Hercules, across Ophiuchus, and up on the eastern side of him to Aquila.

Celestial cartographers have divided the single constellation into two distinct and separated parts called Serpens Caput or Head of the Serpent and Serpens Cauda or Tail of the Serpent. Serpens is a distinct and pretty line of small stars, fairly close together and sometimes almost in pairs, winding around through this region of the skies; and it is particularly well defined in August and September, when it is rather low in the southern heavens.

## CAPRICORNUS

Capricornus, or the Sea Goat, the tenth sign and eleventh constellation of the Zodiac, lies southeast of Aquila. The sun occupies it from January 18th to February 14th. It has no conspicuous stars, and no figure definitely outlined that can be described; and yet on a clear night the comparatively faint stars dancing about in it may suggest a goat, if one does not try to be too definite. Alpha and Beta, which are in the head of the Goat, lie about twenty degrees south of Altair and point towards it. They are about two and one-half degrees apart, and are attractive out of all proportion to their brilliancy. Near Alpha is a faint fifth-magnitude star that is even a better test of vision than Alcor. Capricornus lies east of Sagittarius and south of the Dolphin.



## AQUARIUS

Aquarius covers the part of the Zodiac that the sun occupies from February 14th to March 14th. The best time to observe the constellation is when it is farthest from the sun, in August, September, October, and November. Aquarius has three stars of the third magnitude and none brighter. It is of a peculiar and straggling shape, extending around Capricornus on the north and east. The eastern part of it is south of Pegasus, and the western part is south of Delphinus. It is supposed to represent a man pouring water from an urn. Its distinctive feature to the naked eye is a waving line of faint twinkling stars that somewhat suggest a stream of water, and which lead down to our old acquaintance Fomalhaut, in the mouth of the Southern Fish (*Piscis Austrinus*). A bright Y-shaped asterism marks the water jar. Aquarius is the eleventh sign and the twelfth constellation of the Zodiac.

## PISCIS AUSTRINUS

Of this constellation all we need to know is that it consists of Fomalhaut and the few fifth-magnitude stars surrounding it.

## CETUS

Cetus is a large constellation lying almost entirely south of the equator. It represents a huge sea monster and is sometimes called the Whale. The head, marked by a small pentagon of stars, is on the equator and lies directly below Aries, about twenty-two degrees south of the principal star (Hamal). The body stretches along below Pisces and down to Aquarius, over a distance of more than forty degrees. The body is outlined by a four-sided figure easily distinguishable in this rather sparsely-starred region, and the tail is marked by the only second magnitude star in the constellation—a pretty yellow star sometimes called Diphda. The most noted object in Cetus is the star Mira (The Wonderful), its Greek letter name being Omicron Ceti. It is a remarkable variable star and changes in light all the way from the second to the ninth magnitude. It is bright enough to

be visible about a month or two each year, and reaches a maximum of brightness about every eleven months. However, its maxima are not easily visible each year because some occur at seasons of proximity to the sun. Sometimes it is barely visible and at other maxima may attain the second magnitude. Once, in 1779, it is reported to have been as bright as Aldebaran. It has been under observation for more than three hundred years.

## ORION

Orion is the most brilliant of all the southern constellations. It has been fully described in the chapter on "Orion's Bright Stars." It rises just after Taurus and is visible from late October to May.

## LEPUS

Lepus, or the Hare, is a small constellation just south of Orion. It is very pretty in the winter, but too low down in the sky to be seen clearly except when Orion is well up in the heavens.

## ERIDANUS

Eridanus is a stream of stars beginning just west of the Orion Nebula and meandering west and south among the stars until it disappears below the southern horizon. Achernar, its only first-magnitude star, is too far south for us to see it. Though most of the stars are faint, the constellation is easily traced because so few stars lie near it.

## CANIS MAJOR

Canis Major, or the Great Dog, lies just east of Lepus and south of Monoceros. It can always be identified by its bright star, Sirius. Canis Major represents a dog standing on his hind feet and pawing in the air in front of Lepus, the Hare. Sirius marks the nose of the dog; Adhara, a star nearly as bright as the first magnitude, is in the flank; and Mirzam, or Beta, scarcely less

bright, is in one fore-paw. There are two other second-magnitude stars, making the constellation, as a whole, one of considerable brilliancy, even without the great Sirius. It contains forty-one stars that can be seen without a glass.

## HYDRA

Hydra, the Water Snake, is a straggling constellation that covers between ninety and one hundred degrees in its length, and has but one star of any considerable brightness. This is a second-magnitude star, sometimes known as Cor Hydræ (the Heart of the Snake), sometimes as Alpha Hydræ, and sometimes as Alphard, the Solitary One. It is a reddish star, about twenty degrees south and a little west of Regulus; and, as its name indicates, Alphard has no other bright star very near it.

Cor Hydræ rises about half an hour later than Regulus, reaches the meridian three-quarters of an hour earlier, and sets two and one-half hours before Regulus disappears. The Water Snake itself has its head directly south of Præsepe in Cancer and about half-way between Procyon and Regulus. The body runs southeasterly (outlined by Alphard and numerous faint stars), below Leo and Virgo, and ends under Libra. Except for its one bright star, Hydra is of no particular interest.

The back of Hydra serves as a resting-place for two very pretty constellations known as Corvus the Crow and Crater the Cup. Corvus lies a little to the southeast of Spica, and has four principal stars forming a four-sided figure with a little star out at one side, making the outline somewhat resemble that of Libra. The four stars are rather bright third-magnitude ones, and the constellation is a conspicuous one during the spring and early summer months when Virgo and Leo are in prominent places in the sky.

## PISCES

Pisces is the twelfth sign and first constellation of the Zodiac. It has no bright stars, and is best described as the body of stars occupying the space from Aquarius to Aries and running below Andromeda and Pegasus and north of Cetus.

Pisces is supposed to represent two fishes tied together with ribbons, a very inappropriate combination; and it is not impossible to imagine that the faint stars running through it do outline something of the sort, but certainly not with any definiteness. The reviser of this book maintains that the lines representing the cords or ribbons originally were streams of water sprayed from the mouths of the two fishes and splashing against the back of Cetus.

Pisces should not be confused with *Piscis Austrinus*, or the Southern Fish, which is an entirely different constellation. The most important thing about Pisces is that in it lies the point where the ecliptic, or path of the sun, crosses the equator. This point is passed by the sun on its way north in the spring and is called the vernal equinox. The line running from one pole to the other and passing through the vernal equinox is the one from which astronomers measure the right ascension of the stars (equivalent to longitude on the earth). For this reason Pisces is an interesting constellation, and it is to be regretted that it has no bright star to mark so important a point. However, since the vernal equinox slowly moves, occupying each of the twelve Zodiacal constellations in turn, over an interval of 25,000 years, no star could represent the equinoctial point for very long.

### XXX

## Individuality of the Stars

WE ought now to begin to feel at home in the presence of the general company of stars. We have been introduced to most of the choicer and greater individuals among them and have had a general view of their brilliant assemblage as it appears from night to night and from season to season and a slight insight into their common life. If any permanent relation is to ensue we must now—each for himself—make our own friends among them. What most interests one may not most interest another. We do not feel alike in regard to people or other objects in nature, so why should we in regard to the stars? The more individuality any thing or person has, the surer are we to have a decided feeling of attraction or repulsion for it, a readiness to admit it to our regard or to let it pass with the unnumbered individuals that make no impression upon us.

More than any other natural objects, the stars, as they appear to us in the skies, seem to me to have individuality. Each lovely flower that we see is one of a species, and however much character we may find in the species, every individual is known only by its being a duplicate of thousands of others in petal, sepal, stamen, and anther. The shy hermit-thrust and the friendly robin in their perfect normal state must conform to their kind even to the color of their bills. But there is only one Vega, one Capella, one Arcturus, and they are no more duplicated in nature than they are in name.

What the general character of a star assumes to us depends upon ourselves, the peculiarities to which we are sensitive and the play our fancy takes in the interpretation of them. But the traits that go to make up this general character have their basis in facts and are by no means products of a freakish imagination. Nearly every one has some sensitiveness to color and will

receive an impression when he marks Antares as red, Vega as bluish, Castor as tinged with green, and the various other stars as having their own peculiar tints. This will have an additional meaning, perhaps, when he remembers that these great white stars with tints of various colors represent extremely hot stars. It will then be but natural to remember that Capella, Procyon, Pollux, and Arcturus and the other stars of a like yellow hue are somewhat cooler; while Antares, Betelgeuse, and Aldebaran and similar red stars are the coolest of all. Betelgeuse even allows his eyelids to droop at intervals and thus becomes a variable star, which is thought to be one of the marks of advanced age in star life.

One must also get some impression of individuality as he learns something of the motion and distance and dimensions of a star. Our personal estimate of a star and relation to it cannot but be influenced by such a fact as that Spica and Deneb are twelve and forty times respectively more distant than comparatively near Altair. Or we may learn that one star whirls around its companion in seven hours and another spends hundreds of years accomplishing the same task. One star may fly at the rate of two hundred miles in a second and another loiters along at a pace scores of times slower. One star is plunging towards us and another is receding from us, and whatever its rate or direction, each star is pursuing its own peculiar path.

The length of its stay with us, the time of year that we see it most, and the personal association we have with it must also have some effect on our relation to a star. There must always be less of intimate pleasure in the presence of a star like Fomalhaut, for instance, which comes for only a few months each year and makes a short, swift passage along the southern edge of our skies, than with such stars as Capella and Vega, who turn their shining faces towards us throughout the entire night so many months in the year. There is likely to be more warmth of feeling for the bright Spica, who shines during all the rich soft nights of summer, than for the much brighter Rigel, who dances along almost exactly the same path during the cold, crisp winter nights. Certain stars that may have the intensest charm and companionability for us if we first learn to know them as they hang during the rich black nights under the great dome that arches over spacious fields and valleys, may scarcely attract a

passing thought from us if we meet them first in the narrow alleyways of the city streets, or even in the hazy, half-lighted atmosphere that envelops the higher points in the city.

Even the name of a star imparts some flavor to one's general notion of it. Some names are essentially feminine, some distinctly masculine, some have a meaning that cannot be overlooked, and some appeal to one's fancy by mere association of sound.

As we come into closer acquaintance with the stars and give heed to them, such characterization comes about as naturally as it does with any other objects that play into our lives. In my journal of out-door things I find mention of perhaps thirty or more stars that seem to me to have well-defined traits which are indissolubly attached to my idea of the star. Antares, for example, seems old but important and impressive, Capella fair and mature, Vega young and fresh and joyous, Arcturus full of dignity, Sirius brilliant without much impressiveness, Betelgeuse gloomy even to the point of being foreboding, and suggestive of the sullen redness of the light before whose warning eye the swift electric trains come to a dead stop. Dubhe and Merak seem fair and smiling as they graciously point the way to the North Star, Mizar suggests friendliness, Alcor faithfulness, Aldebaran sturdiness, the Pleiades delicate charm, Fomalhaut serenity, Castor and Pollux gentleness and kindness and affection, Mira instability, and Polaris its opposite—stability and responsibility.

Many more might be added to this list, but I have no desire or expectation that others may have the same impressions that I have concerning individual stars. Circumstances may lead them to lay stress upon quite different traits which are as truly characteristic of the star as the ones that have appealed most to me. These have been mentioned to indicate the pleasant familiarity that one may assume with the individuals composing this great body of nature that lies above our eyes every night of our lives, and to create this sort of an interest in them has been the sole aim of this book.

.





# Index

- ACHERNAR (Alpha Eridani), 11; pronunciation, 11; magnitude, 81.  
 Acrux (Alpha Crucis), 11, 77, 81.  
 Adara (Epsilon Canis Majoris), 136.  
 Albireo (Beta Cygni), a double star, 34.  
 Alcor, companion to Mizar, 77, 106-107.  
 Alcyone, 127.  
 Aldebaran (Alpha Tauri), pronunciation, 11; chapter xi; color, 48; in constellation Taurus, 48; location, 48; meaning of name, 49; winter star, 49; brilliancy, 50; lies in moon's path, 50; "royal" star, 50; when visible, 50; magnitude, 81.  
 Alderamin (Alpha Cephei), 109.  
 Algenib (Gamma Pegasi), in the square of Pegasus, 116.  
 Algol (Beta Persei), 120; known as "The Ghoul" or winking star, 120; magnitude, 120.  
 Alioth (Epsilon Ursæ Majoris), in the Great Dipper, 105.  
 Alkaid (Eta Ursæ Majoris), 105.  
 Almach (Gamma Andromedæ), 120.  
 Alpha Andromedæ (Alpheratz), 119.  
 Alpha Arietis (Hamal), 122.  
 Alpha Canum Venaticorum (Cor Caroli), 111.  
 Alpha Cassiopeia (Schedar), 109.  
 Alpha Centauri (Rigel Kent), 12; pronunciation, 10; double star, 12; nearness to earth, 12; Southern Star, 12; magnitude, 81.  
 Alpha Cephei, 109.  
 Alpha Coronæ Borealis (Alphecca), 90.  
 Alpha Crucis, 11; pronunciation, 11; magnitude, 81.  
 Alpha Draconis as Pole Star, 102.  
 Alpha Hydræ (Cor Hydræ), 137.  
 Alpha Pegasi (Markab), 116.  
 Alpha Persei (Mirfak), in the segment of Perseus, 120.  
 Alpha Ursæ Majoris (Dubhe), 105.  
 Alpha Ursæ Minoris (Polaris), 6, 101.  
 Alphecca (Alpha Coronæ), chief star in Corona Borealis, 112; rising of, 112.  
 Alpheratz (Alpha Andromedæ), 116, 119.  
 Altair (Alpha Aquilæ), 11; pronunciation, 11; chapter viii; how to locate, 35; when visible, 36-39; development, 39; distance from the earth, 39; magnitude, 81.  
 Alwaid (Beta Draconis), 110.  
 Andromeda, 117; the constellation, 117; diagram, 118; famous nebula, 119.  
 Antares (Alpha Scorpii), 2; Southern Star, 7; pronunciation, 11; chapter ix; location, 40; Southern Star, 40; when visible, 40-43; color, 43; in the constellation Scorpius, 43; red giant, 43; binary system, 44; distance from the earth, 50; "royal" star, 50; magnitude, 81.  
 Aquarius (The Water Bearer), 135; related to Pegasus, 116.  
 Aquila (The Eagle), 115; stars of, 115.  
 Archer, the (Sagittarius), 113.  
 Arcturus (Alpha Bootis), splendor, 2; pronunciation, 10; chapter iv; brilliancy, 20; how to locate, 20; rising time, 20-21; color, 21; diameter, red giant, 21; distance from the earth, 22; speed, 22; in the constellation Bootes, 23; magnitude, 81; part of the diamond of Virgo, 131.  
 Argo Navis, 99.  
 Arided, *see* Deneb.  
 Aries (The Ram), 122; first sign of the Zodiac, 122.

- Arrow, the (Sagitta), 115.  
 Asterism, 12, 99.  
 Asterope, 127.  
 Atlas, a double star, 127; one of the brightest stars in the Pleiades, 127.  
 Auriga (The Charioteer), 19; 120-121.  
  
 BARNARD'S STAR, 89.  
 Becrux (Beta Crucis), 11, 77.  
 Bellatrix, in Orion, 55.  
 Benetnasch, *see* Alkaid.  
 Berenice's Hair (Coma Berenices), 90.  
 Beta Andromedæ (Mirach), 119.  
 Beta Aquilæ, 115.  
 Beta Arietis (Sheratan), 122.  
 Beta Aurigæ (Menkalinan), 121.  
 Beta Canis Majoris (Mirzam), 136.  
 Beta Cassiopeïæ (Caph), 108.  
 Beta Centauri, 11; pronunciation, 11; magnitude, 81.  
 Beta Cephei, 109.  
 Beta Ceti (Diphda), 135.  
 Beta Crucis (Becrux), 11; pronunciation, 11; magnitude, 81.  
 Beta Draconis (Alwaid), 110.  
 Beta Leonis (Denebola), 68.  
 Beta Pegasi (Scheat), 116.  
 Beta Persei (Algol), 120.  
 Beta Ursæ Majoris (Merak), 106.  
 Beta Ursæ Minoris (Kochab), 103.  
 Betelgeuse (Alpha Orionis), 3; pronunciation, 11; irregular variable, 52; red giant, 52; distance from the earth, 54; magnitude, 81.  
 Big Dipper, *see* Great Dipper.  
 Binaries, spectroscopic, 97.  
 Binary systems, chapter xx.  
 Bootes, 23; 111-112.  
 Bull, the (Taurus), 48, 123.  
  
 CANCER (The Crab), 129; fourth sign of the Zodiac, 129; Præsepe in, 129.  
 Canes Venatici (The Hunting-Dogs), 111.  
 Canis Major (The Great Dog), 136.  
 Canis Minor (The Smaller Dog), 109.  
 Canopus (Alpha Carinæ), brilliancy, 11; distance from the earth, 11; pronunciation, 11; ran away with one of the Pleiades, 11, 128; Southern Star, 11; magnitude, 81.  
 Capella (Alpha Aurigæ), 3, 19; pronunciation, 10; chapter iii; brilliancy, 17; how to find, 17; rising time, 17-18; when visible, 17-18; color, 18; rarefied body, 18; a double star, 18-19, 92; magnitude, 81.  
 Caph (Beta Cassiopeïæ), 108.  
 Capricornus (The Goat), 134; relation to Auriga, 121; tenth sign of the Zodiac, 134.  
 Carina, (The Keel), 99.  
 Carlyle, Thomas, quoted, 99.  
 Cassiopeia, chapter xxiv.  
 Castor (in Gemini), 57; brilliancy, 57; rising, 57; course, 58; superstition, 58; distance from the earth, 59; a multiple star, 59, 90.  
 Celæno, 127.  
 Cepheus, chapter xxiv.  
 Cetus (The Whale), 135.  
 Clusters, moving, 90.  
 Coma Berenices, 90.  
 Concerning the brightest stars, chapter ii.  
 Constellations, chapters XXI-XXIX; for details refer to constellation name.  
 Cor Caroli (Alpha Canum Venaticorum), 111; part of the diamond of Virgo, 131.  
 Cor Hydræ (Alpha Hydræ), only bright star in Hydra, 137.  
 Corona Borealis (Northern Crown), 112.  
 Corvus (The Crow), 137.  
 Crab Nebula, 88.  
 Crater (The Cup), 137.  
 61 Cygni, 34.  
 Cygnus (The Swan), 34, 114.  
 Cynosura (Polaris), the end of the dog's tail, 102.  
  
 DELPHINUS (The Dolphin, or Job's Coffin), 35, 38, 115.  
 Delta Andromedæ, 119.  
 Delta Cassiopeïæ, 108.  
 Delta Leonis (Zosma), 130.  
 Delta Pegasi (Alpheratz), 116, 119.  
 Delta Ursæ Majoris (Megrez), 105.  
 Deneb (Arided), (Alpha Cygni), head of Northern Cross, 6; pronunciation, 11; chapter vii; rising time, 30; when visible, 30-33; meaning of the name, 32; color, 33; distance from the earth, 33; supergiant, 33; magnitude, 81.  
 Denebola (Beta Leonis), location, 68; meaning of name, 68; part of the diamond of Virgo, 131.  
 Diamond of Virgo, 131.  
 Diphda (Beta Ceti), 135.  
 Distances of the stars, chapter xix.  
 Dog Stars, the two, chapter xiv; *see also* Procyon and Sirius.  
 Dolphin, the (Job's Coffin), *see* Delphinus.  
 Double stars, chapter xx.

Draco (The Dragon), 109-110.  
 Dubhe (Alpha Ursæ Majoris), 105.  
 Dust, interstellar, 71.

EINSTEIN'S THEORY OF RELATIVITY, 71.

Electra, 127.

Emerson, quoted, 4.

Enif (Epsilon Pegasi), 116.

Epsilon Canis Majoris (Adara), 136.

Epsilon Cassiopeia, 108.

Epsilon Lyra, a double-double star, 91.

Epsilon Pegasi (Enif), 116.

Epsilon Ursæ Majoris (Alioth), 105.

Eridanus (The River), 136.

Eta Aquilæ, variable star, 115.

Eta Cassiopeia, double star, 108.

Eta Ursæ Majoris (Alkaid), 105.

Etamin (Gamma Draconis), 110.

FOMALHAUT (Alpha Piscis Austrini),  
 when visible, 2; Southern Star, 7;  
 pronunciation, 11; chapter x; bril-  
 liancy, 45; location, 45; rising time,  
 46; "royal" star, 46; nearness to  
 earth, 47; magnitude, 81.

GAMMA ANDROMEDE (Almach), 119.

Gamma Aquilæ, 115.

Gamma Cassiopeia, 108.

Gamma Draconis (Etamin), 110.

Gamma Pegasi (Algenib), 116.

Gamma Ursæ Majoris (Phecda), 106.

Gamma Ursæ Minoris, 103.

Gemini (The Heavenly Twins),  
 chapter XIII, 129; *see also* Castor  
 and Pollux.

Goat (Capricornus), relation to  
 Auriga, 121.

Gravitation, 95.

Great Bear (Ursa Major), 12-15;  
 the stars of, chapter XIII; always  
 visible, 104; its stars form the  
 Great Dipper, 104.

Great Dipper, the 6, 105; *see also*  
 Great Bear.

Great Dog (Canis Major), the, 136.

Guard stars, 102-103.

HADAR (Beta Centauri), 11.

Hamal (Alpha Arietis), chief star  
 in Aries, 122.

Heavenly Twins, the (Gemini),  
 chapter XIII.

Harvard Photometry, 79.

Hercules, 110, 113; diagram, 113.

Hunting-Dogs, the (Canes Venatici),  
 111.

Hyades, the, 48-49, 90.

Hydra (The Water Snake), 137.

INDIVIDUALITY OF THE STARS, chapter  
 XXX.

Job's Coffin, *see* Dolphinus.

Jupiter, the planet, 15-16.

KAPPA CASSIOPEIA, 108.

Kappa Orionis (Saiph), 55.

Kochab (Beta Ursæ Minoris), 103.

LATITUDE, 8.

Leo (The Lion), 130; Denebola, 130;  
 fifth sign of Zodiac, 130; Regulus,  
 130.

Leonids, 69.

Lepus (The Hare), 136.

Libra (The Scales), 131.

Light, of the stars, chapter XVIII;  
 stellar magnitude, 78-83; speed of  
 light, 87.

Little Bear, the (Ursa Minor), chap-  
 ter XXII; north pole within its  
 boundaries, 102; Polaris, 102.

Little Dipper (Ursa Minor). *see*  
 Little Bear.

Little Dog, *see* Canis Minor.

Lost Pleiad, the, 127.

Lyra (The Lyre), 29, 114.

MAGNITUDE OF THE STARS, 72.

Maia, 127.

Markab (Alpha Pegasi), 116.

Mars, the planet, 15-16.

Megrez (Delta Ursæ Majoris), 106.

Menkalinan (Beta Aurigæ), 121.

Merak (Beta Ursæ Majoris), 106.

Mercury, the planet, 15.

Merope, 127.

Meteors, 69.

Milk Dipper, the, in Sagittarius, 133.

Milky Way, 71.

Mira, the Wonderful (Omicron  
 Ceti), 76.

Mirach (Beta Andromedæ), 119.

Mirfak (Alpha Persei), 120.

Mirzam (Beta Canis Majoris), 136.

Mizar (Zeta Ursæ Majoris), 77, 106;  
 first double star discovered, 94;  
 quadruple star, 96.

Monoceros (The Unicorn), 136.

Moon, apparent diameter of, 125.

Multiple stars, 97.

NAMES OF THE STARS, chapter XVII.

Nebula, Orion, 55.

Neptune, the god, associated with  
 Auriga, 121.

Neptune, the planet, 15.

Newton, Sir Isaac, 95.

North Star, the (Polaris), 6, 101.

Northern Cross, the (Cygnus), 6.

Northern Crown (*Corona Borealis*), 112.

Number of stars, chapter xvi.

OMICRON CETI (*Mira*, the Wonderful), 135-136.

Ophiuchus and Serpens, 134; location, 134; when visible, 134.

Orion, and a frosty night, 2; chapter xii; bright stars in, 52; mythology, 52; sword of, 52; belt of, 55; most admired, 55; nebula, 55; "trapezium," 55; when visible, 56; three kings, 136.

PALOMAR OBSERVATORY, 73.

Parallax of a star, finding the, 88.

Pegasus (*The Winged Horse*), 116-117.

Perseus, 119; segment of, 120.

Phecda (*Gamma Ursæ Majoris*), 106.

Photometer, 79.

Pisces (*The Fishes*), 137.

Piscis Austrinus (*Southern Fish*), 135.

Planets, the, 15; colors of, 16.

Pleiades, the, 49; lost, 11, 127; moving cluster, 90; chapter xxvii; influence of, 124; location of, 124; principal stars in, 125; brightest stars of, 126; distance from the earth, 126; enwrapped in cosmical fog, 126; time of rising, 126; number of stars in, 127.

Pleione, the lost Pleiad, magnitude, 127.

Pluto, the planet, 15.

Polaris (*Alpha Ursæ Minoris*), 6, 101.

Pole Star, *see* Polaris.

Pollux, pronunciation, 11; in constellation Gemini, 57; winter star, 57; superstition, 58; distance from the earth, 59.

Præsepe, a star cluster in Cancer, 90.

Procyon, pronunciation, 11; a winter star, 63; in *Canis Minor*, 64; meaning of name, 64; the smaller Dog Star, 64; a double star, 65; nearness to earth, 65; magnitude, 81.

Proper motion of a star, 92.

Proxima (*Alpha Centauri*), 12.

Puppis, the Stern, 99.

Pyxis, the Compass, 99.

RADIO STARS, 88.

Red giants, nature of, 22, 43.

Regulus (*Alpha Leonis*), pronunciation, 11; chapter xv; location, 66; brilliancy, 67; when visible, 67;

distance from the earth, 68; in constellation Leo, 68; meaning of name, 68; magnitude, 81.

Rigel (*Beta Orionis*), a southern star, 7; pronunciation, 11; brilliancy, 54; color, 54; distance from the earth, 54; location, 54; rising, 54; a winter star, 54; magnitude, 81.

Rigel Kent (*Alpha Centauri*), 10, 12.

Ring Nebula, the, 114.

Rising and setting of the stars, the, chapter i.

River, the (*Eridanus*), 136.

"Royal" stars, 50.

SAGITTA (*The Arrow*), 115.

Sagittarius (*The Archer*), 113; ninth sign of the Zodiac, 113.

Saiph (*Kappa Orionis*), 55.

Saturn, the planet, 15-16.

Scheat (*Beta Pegasi*), 116.

Schedar (*Alpha Cassiopeæ*), 109.

Schmidt camera, 73.

Scorpius, 132; eighth sign of the Zodiac, 132.

Segment of Perseus, 120.

Serpens (*The Serpent*), 134; *see also* Ophiuchus.

Sheratan (*Beta Arietis*), 122.

Sickle, the (*Leo*), 130.

Sirius (*Alpha Canis Majoris*), departure of, 7; location, 7; southern star, 7; pronunciation, 10; companion of, 22, 44, 64; brilliancy, 61; Dog Days, 61; Dog Star, 61; in *Canis Major*, 61; a winter star, 61; course, 62; Orion's Dog, 62; rising, 62; when visible, 62; color 63; distance from the earth, 63; magnitude, 81.

Sound, speed of, 87.

Southern Fish, *see* *Piscis Austrinus*.

Southern stars, 7; chapter xxix.

Speed of light, 87.

Spica (*Alpha Virginis*), southern star, 7; pronunciation, 11; chapter v; brilliancy, 24; how to find, 24; color, 25; distance from the earth, 25; hour of setting, 25; meaning of name, 25; when visible, 25; magnitude, 81; part of the diamond of Virgo, 131.

Square of Pegasus, 116.

Stars, the, time of rising and setting, 5, 8; chapter ii; pronunciation, 10-11; colors of, 62-63; magnitudes, 81; binary systems, chapter xx; multiple, 97; individuality of, chapter xxx.

Sun, the, distance from the earth, 87.  
Sword of Orion, 55.

TAURUS (The Bull), 48, 123; the Hyades in, 123; and Orion, 123; the Pleiades in, 123; second sign of the Zodiac, 123; stars in, 123.

Taygeta, 127.

Theta Leonis, 130.

Theta Orionis, double star, 94, 96.

"Three Kings," the, 55.

Thuban (Alpha Draconis), formerly the Pole Star, 102.

Trapezium in Orion, *see* Theta Orionis.

Triangulum, stars in, 122.

Twins, Heavenly (Gemini), chapter XIII.

Two Dog Stars, the, chapter XIV; *see also* Procyon and Sirius.

URANUS, the planet, 15.

Ursa Major, *see* Great Bear.

Ursa Minor, *see* Little Bear.

VEGA (Alpha Lyrae), harbinger of spring, 1, 2; rising, 6, 27-29; pronunciation, 10; chapter VI; compared with Arcturus, 26; brilliancy, 27; how to find, 27; in constellation Lyra, 29; when visible, 29; magnitude, 81.

Vela (The Sail), 99.

Venus, the planet, 15-16.

Virgo (The Virgin), 131; sixth sign of the Zodiac, 131.

Virgo, Diamond of, 131.

WAIN, 107.

White dwarfs, 22, 44, 64, 65.

Winged Horse, *see* Pegasus.

ZETA URSAE MAJORIS (Mizar), 77.

Zosma (Delta Leonis), 130.

